



Estimation of Breath Holding Time (BHT) among Dental Students

Aashiq Mohamed^a, R. Gayatri Devi^{b*} and A. Jothi Priya^b

^a Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Science (SIMATS), Saveetha University, Chennai-600077, India.

^b Department of Physiology, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Science (SIMATS), Saveetha University, Chennai-600077 India.

Authors' contributions

This work was carried out in collaboration among all authors. Author AM Literature search, data collection analysis, manuscript drafting. Author RGD Data Verification, Manuscript drafting. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i58A34108

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/78933>

Original Research Article

Received 06 October 2021

Accepted 13 December 2021

Published 15 December 2021

ABSTRACT

Background: Breath holding time is the time taken by an individual to hold his/her breath as long as they can. During voluntary breath holding, tissues continue to utilize oxygen and liberate carbon dioxide. Therefore, during breath holding arterial pO₂ falls and pCO₂ rises. Normal Breath holding time (BHT) is 45-55 seconds which is estimated at 2014. The main aim of this study is to assess the breath holding time among healthy dental students.

Materials and Methods: A minimal number of sample sizes have been taken into account with regards to their BMI. The sample size was divided into two groups as gender comparison. Students were detail explained about the procedure and breath holding time has been measured. The statistical analysis was done by ANOVA test in SPSS Software-23 and an independent t-test was calculated.

Results: BHT was found to be significantly low among females compared to males. A statistically significant negative correlation was observed when BHT is correlated with gender because BHT differs in both the gender.

Conclusion: The present study revealed that breath holding time is less in females compared to males. Therefore, males are healthier than females.

[≡] Assistant Professor;

*Corresponding author: E-mail: gayatridevi@saveetha.com;

Keywords: *Breath holding time; gender comparison; BMI; Oxygen; Carbon Dioxide; Innovation techniques.*

1. INTRODUCTION

Breath holding time is the time taken by an individual to hold his/her breath as long as they can [1]. Breath holding involves the tissues which use oxygen and then produce carbon dioxide [2]. Breath holding increases carbon dioxide in the arterial region resulting in vasodilation of resistance arterioles which increases cerebral blood flow [3]. Evaluation of breath holding time has a significant increase in sensitivity of peripheral chemoreflex, which was noted regarding arterial hypertension, chronic heart failure, etc [4].

Respiratory motion is an abnormal rate of breath due to which causes chest pain and brain tumor [5]. Breath holding for short intervals is a widely applied method to reduce respiratory motion [6]. In standard clinical practice, patients are instructed to hold their breath [7]. In addition to Positron Emission Tomography and Computed Tomography (PET-CT) Scans and radiotherapy have been used in an MR setting recently [8]. Breath hold challenge is used in the clinical setting for assessment of cerebrovascular reactivity [9].

Hypoxia is a condition in which a certain body part suffers from a decreased rate of partial pressure of oxygen levels whereas Hypercapnia is an increased rate of partial pressure of carbon dioxide levels [10]. Hypercapnia is commonly measured as a surrogate for the increased arterial carbon dioxide levels [11]. Hypoxia of oxygen were seldom used to account for Cerebral blood flow(CBF) changes during breath holding due to the belief that vasodilatory effect of increased artery partial pressure of carbon dioxide dominates that of decreased arterial partial pressure in oxygen [12]. The role of both hypoxia and hypercapnia in breath holding deserves to be examined because arterial PCO_2 and PO_2 have been reported to work synergistically to interact with chemoreceptors [13].

Some studies described the potent vasodilatory nature of arterial carbon dioxide in humans [14] which can be manipulated using inhalation of air with increased carbon dioxide [15] [16]. The

present study mainly stimulates individuals' behaviour in healthy volunteers. This allows more flexibility in the experiment duration and removes the confounds of age-related changes in order to isolate the effects of poor breath hold performance on the Blood Oxygen Level Dependent (BOLD) measurements [17]. Coronavirus is being viral all over the world, this study has the main role to compare breath holding time between males and females to check whether they have proper lung functioning so that they don't get affected by any external parameters [18] [19]. The main aim of this study is to explore the gender comparison of Breath Holding Time among dental students.

2. MATERIALS AND METHODS

A cross-sectional study was conducted in a private dental college among first years with a sample size of 100 students. The participants are selected randomly. Breath holding time was recorded among each individual by allotting them a nose piece after them holding their breath. The breath holding duration and the measurement of distress tolerance which denotes the maximum experienced distress and the participants were instructed to hold their breath as long as they can. To calculate breath holding timings, participants were instructed to hold their breath as long as they could. To calculate BHTs, participants were assessed to record the starting time from the time they held their breath. The time is calculated using a stopwatch and the readings of BMI are collected by calculating the individual's height and weight. Data was collected and analysed using SPSS software version 23 and independent t-test was calculated.

3. RESULTS

From the present study, BMI mean value in males is 23.1086 ± 4.6270 and in females is 22.5981 ± 4.3723 (Fig. 1). BHT mean value in males is 42.17 ± 39.64 and in females is 26.96 ± 9.435 (Table 1). Since male population has a high mean value, it is evident that male have more breath holding time than females under this sample size (Fig. 2).

Table 1. Mean SD and SE of BMI and BHT. females have participated in this survey rather than males. the mean, standard deviation and standard error mean values are more for males than females

| | Groups | N | Mean | Std. Deviation | Std. Error |
|-----|--------|----|---------|----------------|------------|
| BMI | Male | 29 | 23.1086 | 4.62700 | .85921 |
| | Female | 70 | 22.5981 | 4.37237 | .52260 |
| BHT | Male | 29 | 42.17 | 39.640 | 7.361 |
| | Female | 70 | 26.96 | 9.435 | 1.128 |




Fig. 1. The bar graph represents the mean value of BMI of both genders. X axis represents gender and Y axis represents mean values of BMI of the participants. Green denotes males and Red denotes females. males have more BMI than females




Fig. 2. The bar graph represents the mean value of BHP of both genders. X axis represents gender and Y axis represents mean values of BHP of the participants. Green denotes males and Red denotes females. Males have more BHP than Females

4. DISCUSSION

Breath holding is a safe and reliable method for assessing the sensitivity of peripheral chemoreflexes to carbon dioxide in patients with heart failure [20]. A normal breath holding capacity in an untrained individual is 30 sec to 2 min [21]. In this study, the maximum limit of breath holding capacity of a person was very challenging to measure [22]. Therefore by understanding how these mechanisms interact to determine the break point of a breath hold which remains a challenging area of integrative physiology [23]. By comparing the previous studies which are done similar to the present study [1], the breath holding spells of pediatrics have more significance towards breath holding capacity in children [24]. Then the evaluation of breath holding tests for peripheral chemoreflex sensitivity in patients was performed and also about the chronic heart failure [25].

The study correlated well with breath holding with post bronchodilator(26,27), forced expiratory volume (FEV)[28], forced vital capacity (FVC)[29], peak expiratory flow rate (PEFR) is normal which was similar to the findings observed in the current study [25,30]. Mariappan *et al* conducted a study on the effect of physical training among Indian people (31), it enhances the respiratory efficiency by increasing the strength of diaphragm and intercostal muscles which were similar to the findings that male have more breath holding time than females in this study [32]. It also increases vital capacity and prolongs the breath holding time [33]. Then by comparing the impact of linear vestibular stimulation in respiratory endurance which is a vestibular stimulation by the participants to swing from back to front direction [26] [34]. The consensus faced by the overall previous literature is by the impact of peripheral chemoreflex related to chronic heart failure, obesity, etc [25] [35].

Based on these findings, conclude that males have more breath holding time than females. The number of alveoli per unit area and alveolar volume do not differ between males and females, but it is scientifically proven that males have larger lungs and thoracic cages than females.

5. CONCLUSION

This study was conducted only for normal students in order to check whether they can with stand covid-19. This study clearly shows that

males are higher value than female students since they have more breath holding capacity and further studies should happen on a large scale for better results in the future.

6. LIMITATIONS

This study has less sample size. But this research can be done on a wide scale with a larger sample size and with different age groups and populations due to the fact that this study was only conducted among dental practitioners so that the values of the tests can differ and also awareness of breath holding can easily spread throughout the public.

CONSENT

As per international standard or university standard, respondents' written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

ACKNOWLEDGEMENTS

The author would like to thank Saveetha dental college for the valuable support to conduct the study.

SOURCE OF FUNDING

The present project was funded by

- Saveetha Institute of Medical and Technical Sciences
- Saveetha Dental College and Hospital
- Saveetha University
- NRV Hospital Chennai.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Sridharan G, Ramani P, Patankar S, Vijayaraghavan R. Evaluation of salivary metabolomics in oral leukoplakia and oral squamous cell carcinoma. *J Oral Pathol Med.* 2019 Apr;48(4):299–306.
2. Barabadi H, Mojab F, Vahidi H, Marashi B, Talank N, Hosseini O, et al. Green

- synthesis, characterization, antibacterial and biofilm inhibitory activity of silver nanoparticles compared to commercial silver nanoparticles [Internet]. *Inorganic Chemistry Communications*. 2021;129:108647. Available:<http://dx.doi.org/10.1016/j.inoche.2021.108647>
3. Kwong KK, Belliveau JW, Chesler DA, Goldberg IE, Weisskoff RM, Poncelet BP, et al. Dynamic magnetic resonance imaging of human brain activity during primary sensory stimulation. *Proc Natl Acad Sci U S A*. 1992 Jun 15;89(12):5675–9.
 4. Trembach N, Zabolotskikh I. Breath-holding test in evaluation of peripheral chemoreflex sensitivity in healthy subjects [Internet]. *Respiratory Physiology & Neurobiology*. 2017;235:79–82. Available:<http://dx.doi.org/10.1016/j.resp.2016.10.005>
 5. Bharath B, Perinbam K, Devanesan S, AISalhi MS, Saravanan M. Evaluation of the anticancer potential of Hexadecanoic acid from brown algae *Turbinaria ornata* on HT–29 colon cancer cells [Internet]. *Journal of Molecular Structure*. 2021;1235:130229. Available:<http://dx.doi.org/10.1016/j.molstruc.2021.130229>
 6. Clarizia G, Bernardo P. Diverse Applications of Organic-Inorganic Nanocomposites: Emerging Research and Opportunities. *Emerging Research and Opportunities*. IGI Global; 2019: 237.
 7. Nehmeh SA, Erdi YE, Pan T, Pevsner A, Rosenzweig KE, Yorke E, et al. Four-dimensional (4D) PET/CT imaging of the thorax. *Med Phys*. 2004 Dec;31(12):3179–86.
 8. Kaza E, Symonds-Tayler R, Collins DJ, McDonald F, McNair HA, Scurr E, et al. First MRI application of an active breathing coordinator. *Phys Med Biol*. 2015 Feb 21;60(4):1681–96.
 9. Markus HS, Bunker CB, Kouris K, Costa DC, Harrison MJ. rCBF abnormalities detected, and sequentially followed, by SPECT in neuro-Behçet's syndrome with normal CT and MRI imaging. *J Neurol*. 1992 Aug;239(7):363–6.
 10. Egbuna C, Mishra AP, Goyal MR. Preparation of Phytopharmaceuticals for the Management of Disorders: The Development of Nutraceuticals and Traditional Medicine. Academic Press; 2020:574.
 11. Bright MG, Murphy K. Reliable quantification of BOLD fMRI cerebrovascular reactivity despite poor breath-hold performance. *Neuroimage*. 2013 Dec;83:559–68.
 12. Ezhilarasan D. Critical role of estrogen in the progression of chronic liver diseases. *Hepatobiliary Pancreat Dis Int*. 2020 Oct;19(5):429–34.
 13. Biscoe TJ, Purves MJ, Sampson SR. The frequency of nerve impulses in single carotid body chemoreceptor afferent fibres recorded in vivo with intact circulation [Internet]. *The Journal of Physiology*. 1970;208:121–31. Available:<http://dx.doi.org/10.1113/jphysiol.1970.sp009109>
 14. Kety SS, Schmidt CF. The effects of altered arterial tensions of carbon dioxide and oxygen on cerebral blood flow and cerebral oxygen consumption of normal young men 1 [Internet]. *Journal of Clinical Investigation*. 1948;27:484–92. Available:<http://dx.doi.org/10.1172/jci101995>
 15. Gowhari Shabgah A, Ezzatifar F, Aravindhan S, Olegovna Zekiy A, Ahmadi M, Gheibihayat SM, et al. Shedding more light on the role of Midkine in hepatocellular carcinoma: New perspectives on diagnosis and therapy. *IUBMB Life*. 2021 Apr;73(4):659–69.
 16. J PC, Marimuthu T, C K, Devadoss P, Kumar SM. Prevalence and measurement of anterior loop of the mandibular canal using CBCT: A cross sectional study. *Clin Implant Dent Relat Res*. 2018 Aug;20(4):531–4.
 17. Murphy K, Harris AD, Wise RG. Robustly measuring vascular reactivity differences with breath-hold: normalising stimulus-evoked and resting state BOLD fMRI data. *Neuroimage*. 2011 Jan 1;54(1):369–79.
 18. Kamath SM, Manjunath Kamath S, Jaison D, Rao SK, Sridhar K, Kasthuri N, et al. In vitro augmentation of chondrogenesis by Epigallocatechin gallate in primary Human chondrocytes - Sustained release model for cartilage regeneration [Internet]. *Journal of Drug Delivery Science and Technology*. 2020;60:101992. Available:<http://dx.doi.org/10.1016/j.jddst.2020.101992>
 19. Mudigonda SK, Murugan S, Velavan K, Thulasiraman S, Krishna Kumar Raja VB.

- Non-suturing microvascular anastomosis in maxillofacial reconstruction- a comparative study. *J Craniomaxillofac Surg.* 2020 Jun;48(6):599–606.
20. Nambi G, Kamal W, Es S, Joshi S, Trivedi P. Spinal manipulation plus laser therapy versus laser therapy alone in the treatment of chronic non-specific low back pain: a randomized controlled study. *Eur J Phys Rehabil Med.* 2018 Dec;54(6):880–9.
 21. Skow RJ, Day TA, Fuller JE, Bruce CD, Steinback CD. The ins and outs of breath holding: simple demonstrations of complex respiratory physiology. *Adv Physiol Educ.* 2015 Sep;39(3):223–31.
 22. Prakash AKS, Devaraj E. Cytotoxic potentials of *S. cumini* methanolic seed kernel extract in human hepatoma HepG2 cells [Internet]. *Environmental Toxicology.* 2019;34:1313–9. Available:<http://dx.doi.org/10.1002/tox.22832>
 23. Rajakumari R, Volova T, Oluwafemi OS, Rajesh Kumar S, Thomas S, Kalarikkal N. Grape seed extract-soluplus dispersion and its antioxidant activity. *Drug Dev Ind Pharm.* 2020 Aug;46(8):1219–29.
 24. Inc. KN, Kernel Networks Inc. Docosahexaenoic Acid on Breath Holding Spells in Children [Internet]. *Case Medical Research;* 2019. Available:<http://dx.doi.org/10.31525/ct1-nct03965988>
 25. Trembach N, Zabolotskikh I. Evaluation of Breath-Holding Test in Assessment of Peripheral Chemoreflex Sensitivity in Patients with Chronic Heart Failure. *Open Respir Med J.* 2017 Dec 27;11:67–74.
 26. Santhakumar P, Roy A, Mohanraj KG, Jayaraman S, Durairaj R. Ethanolic Extract of *Capparis decidua* Fruit Ameliorates Methotrexate-Induced Hepatotoxicity by Activating Nrf2/HO-1 and PPAR γ Mediated Pathways [Internet]. *Indian Journal of Pharmaceutical Education and Research.* 2021;55:s265–74. Available:<http://dx.doi.org/10.5530/ijper.55.1s.59>
 27. Saraswathi I, Saikarthik J, Senthil Kumar K, Srinivasan KM, Ardhanaari M, Gunapriya R. Impact of COVID-19 outbreak on the mental health status of undergraduate medical students in a COVID-19 treating medical college: a prospective longitudinal study [Internet]. *PeerJ.* 2020;8:e10164. Available:<http://dx.doi.org/10.7717/peerj.10164>
 28. Tahmasebi S, Qasim MT, Krivenkova MV, Zekiy AO, Thangavelu L, Aravindhan S, et al. The effects of oxygen-ozone therapy on regulatory T-cell responses in multiple sclerosis patients. *Cell Biol Int.* 2021 Jul;45(7):1498–509.
 29. Vivekanandhan K, Shanmugam P, Barabadi H, Arumugam V, Daniel Raj Daniel Paul Raj D, Sivasubramanian M, et al. Emerging Therapeutic Approaches to Combat COVID-19: Present Status and Future Perspectives. *Front Mol Biosci.* 2021 Mar 8;8:604447.
 30. Itoh T, Sato T, Akamatsu T, Shin W. Breath analysis using a spirometer and volatile organic compound sensor on driving simulator [Internet]. *Journal of Breath Research.* 2019;14:016003. Available: <http://dx.doi.org/10.1088/1752-7163/ab30ee>
 31. Wadhwa R, Paudel KR, Chin LH, Hon CM, Madheswaran T, Gupta G, et al. Anti-inflammatory and anticancer activities of Naringenin-loaded liquid crystalline nanoparticles in vitro. *J Food Biochem.* 2021 Jan;45(1):e13572.
 32. Mariappan R, Thomas A, Moorthy R, Raju K, Lakshmanan J, Joy M. Measurement of non-invasive blood pressure in lateral decubitus position under general anaesthesia — Which arm gives more accurate BP in relation to invasive BP - dependent or non-dependent arm? [Internet]. *Indian Journal of Anaesthesia.* 2020;64:631. Available: http://dx.doi.org/10.4103/ija.ija_125_20
 33. R H, Hannah R, Ramani P, Ramanathan A, Jancy MR, Gheena S, et al. CYP2 C9 polymorphism among patients with oral squamous cell carcinoma and its role in altering the metabolism of benzo[a]pyrene [Internet]. *Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology.* 2020;130:306–12. Available from: <http://dx.doi.org/10.1016/j.ooolo.2020.06.021>
 34. Teenumol MJ, Sailesh KS, Johny M, Archana R, Mukkadan JK. Impact of linear vestibular stimulation on respiratory endurance [Internet]. *Research Journal of Pharmacy and Technology.* 2017;10:1691. Available:<http://dx.doi.org/10.5958/0974-360x.2017.00299.2>

35. Wahab PUA, Madhulaxmi M, Senthilnathan P, Muthusekhar MR, Vohra Y, Abhinav RP. Scalpel Versus Diathermy in Wound Healing After Mucosal Incisions: A Split-Mouth Study. J Oral Maxillofac Surg. 2018 Jun;76(6):1160–4.

© 2021 Mohamed et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/78933>