



Profile of Blood Donor Deferrals in a Tertiary Care Centre-Our Institutional Experience

D. Rajeswari Thivya^{a*}, R. Vijayashree^{a^o} and K. Meghanath^{a#}

^a Department of Pathology, Chettinad Hospital and Research Institute, Chettinad Academy of Research and Education, Rajiv Gandhi Salai, Kelambakkam, Tamil Nadu, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i55B33848

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/76668>

Original Research Article

Received 04 October 2021
Accepted 10 December 2021
Published 13 December 2021

ABSTRACT

Background: It is the prime duty of transfusion services to provide safe, adequate and timely need of blood and the blood products. Understanding the reasons for donor deferral can help in planning more efficient recruitment strategies and educate and motivate temporarily deferred donors in order to maintain a safe and adequate supply of blood products.

Aims of the Study: To evaluate and analyze the blood donor deferral pattern in a tertiary care hospital blood bank and to review its influence on blood safety.

Methodology: This retrospective study was conducted in the blood bank, CHRI from the year January 2015 to December 2018. Data like demographic data, clinical history, physical examination, haematological examination, stored in the blood bank was retrieved. The donors will be deferred based on standard WHO guidelines. The collected deferral data was analyzed using SPSS software 2011 version 20.

Results: During the study period there were 7010 registered blood donors. The deferral rate was 5.19%. Among the donor deferrals, females were more commonly deferred ie 31.66%. The deferral rate among voluntary and replacement donors are 4.71% and 11.62% respectively. The rate of permanent deferral (17.86%) was less compared to temporary deferral (82.14%). Among temporary deferral anaemia is the most common cause (27.75%). Seropositive for Hepatitis B is the most common cause for permanent deferral (52.30%).

[≡] Associate Professor;

^o Professor/ HOD;

[#] Assistant Professor;

*Corresponding author: E-mail: drthivya@gmail.com;

Conclusion: In our study temporary deferral is higher this necessities the need of education, motivation of these donors for future donation to maintain a healthy and safe donor pool.

Keywords: Blood donors; deferral; temporary; permanent.

1. INTRODUCTION

The National AIDS Control Organization's (NACO) figures reveal that in India approximately 7.4 million units are donated annually, but the requirement is more than 10 million units [1]. This portrays the blood requirement in our country and more important is to have a safe supply of blood at a reasonable cost. For this purpose it is essential to maintain a healthy pool of voluntary donors. WHO Global Database reveals approximately 1.6 million units were redundant due to variable reasons. Not only this around 13 million blood donors were deferred due to high risk behavior, anemia or a preexisting medical disease [2]. Of the 88.2 million blood donations, 83.3% were donated by voluntary non-remunerated donors, 16.4% by replacement donors, and only by 0.3% as paid donors [3]. The deferral rates varied extensively among countries, ranging from less than 1% to 37% and the median rate of deferral was 12% [3]. Blood donor selection procedure is the cornerstone for safety transfusion practices it perceives to safeguard of the health of not only the recipients but also the donors [4]. All the deferred donors are anticipated to be treated with due respect, care and confidentiality. They should be given a proper and crystal clear explanation for the deferral reason and their doubts should be cleared appropriately. By doing this we are educating and motivating the temporarily deferred donors so that they donate in future.

1.1 Aims and Objective

To evaluate and analyze the blood donor deferral pattern and its causes among blood donors in a tertiary care hospital blood bank and to review its influence on blood safety.

2. MATERIALS AND METHODS

The present study was a retrospective study conducted in the blood bank of Chettinad hospital & research institute. All the recorded data about the blood donors from January 2015-December 2018 for a period of 3 years were retrieved. Details about donors like demographic

data, clinical history, physical examination, hematological examination, vitals were retrieved from the data stored in the blood bank. Institutional ethical clearance was obtained before the commencement of the study (745/IHEC/12-19). In our institute, we follow WHO guidelines for donor deferral. Blood donors were grouped as either fit or unfit donors. Predonation and postdonation deferral data were collected and analyzed. Predonation deferral was based on a standard questionnaire and physical examination including weight, pulse rate, blood pressure, and measurement of hemoglobin concentration. In our blood bank, hemoglobin estimation was by specific gravity method. In terms of blood pressure systolic pressure ranging from 100-140mm of Hg and diastolic pressure ranging from 60-90mm of Hg were considered fit for blood donation. Postdonation deferral was based on screening testing of blood for TTI. ELISA was done for HIV, HBV and HCV. VDRL was done for testing Syphilis. Card test was done for Malaria. The donors were explained the process of blood donation by our staff clearly and adequately before the donation process. The statistics were done using SPSS software 2011 version 20. The values were expressed as frequency and percentage

3. RESULTS

During the 3 year study period there were 7010 registered blood donors. Among them 6701 (95.59%) donated blood. Predonation deferral was 309 (4.41%) and postdonation deferral was 55. Totally 364 were deferred (5.19%) [Table1]. Male donors were more common ie 99.13%. Voluntary donors were more common accounting for 89.59% [Table2]. Among the donor deferrals, females were more commonly deferred in our study ie 31.66%. The deferral rate among voluntary and replacement donors is 4.71% and 11.62% respectively [Table3]. The rate of permanent deferral (17.86%) was less compared to temporary deferral (82.14%). Among temporary deferral Anemia is the most common cause (27.75%) [Table5]. Seropositive for Hepatitis B is the most common cause for permanent deferral (52.30%) [Table 4].

Table 1. Audit of blood donors for 3 years

Year	Number of registered donors	Number donated	Number of deferral	Number of post donation deferral
2017	2294	2181 (95.07%)	113 (4.93%)	14
2018	2346	2246 (95.74%)	100 (4.26%)	24
2019	2370	2274(95.94%)	96(4.06%)	17
Total	7010	6701	309	55

Table 2. Frequency of registered donors

Year	Male	Female	Voluntary	Replacement
2017	2167 (99.35%)	14(0.65%)	1939 (88.90%)	242 (11.10%)
2018	2227 (99.15%)	19(0.85%)	2034(90.56%)	212(9.44%)
2019	2247 (98.81%)	27(1.19%)	2031(89.31%)	243(10.69%)
Total	6641	60	6004	697

Table3. Frequency of deferral donors

Year	Frequency of donor deferral	Male	Female	Voluntary	Replacement
2017	127 (5.53%)	116(91.34%)	11	98 (77.17%)	29
2018	124 (5.28%)	121(97.58%)	3	95(76.61%)	29
2019	113 (4.76%)	108(95.58%)	5	90(79.65%)	23
Total	364	345	19	283	81

Table 4. Causes for permanent deferral

Causes for permanent deferral	2017	2018	2019	Total
Hbs ag	8	15	11	34(52.31%)
HCV	4	8	5	17(26.15%)
HIV	2	1	0	3(4.62%)
VDRL	0	0	1	1(1.54%)
Medications	1	6	2	9(13.85%)
Hemo dialysis	0	0	1	1(1.54%)
	15	30	20	65

4. DISCUSSION

Blood transfusion service is an essential part of the health care system. As per WHO, the minimum requirement to meet a nation's essential blood requirements is taken generally as donation of blood by 1% of its population. But usually the requirements are much higher in countries which have advanced health care systems [5]. According to Siromani et al our nation's blood requirement is calculated as 8.5 to 10 million units/year, but the supply is only 7.4 million units/year [6]. Individuals who approach blood banks intending to donate blood believe them as fit and healthy donors but it is the sole duty of the blood bank to screen them completely and if needed to defer them either temporarily or permanently. This not only aids us to maintain a healthy and safe donor pool but also to safeguard the health of both donors and

recipients. To sustain safe, adequate blood at a reasonable cost we have to educate and motivate temporarily deferred donors for future blood donation. The present study is a retrospective study conducted in the blood bank of Chettinad hospital & research institute. All the recorded data about the blood donation from January 2015- December 2018 for a period of 3 years were retrieved. During the 3 year study period there were 7010 registered blood donors. Among them 6701 (95.59%) donated blood. The rate of deferral in our study is 5.19% and it is lower than most other studies. Most of the Indian and international studies have reported deferral rate ranging from 5% to 24% [7]. In Tufail et al (Pakistan) study the rate is much higher 13.58% [8]. In Henshaw et al (Nigeria) study the deferral rate is 8.69% [9]. Laila et al in their study (Dubai) had a higher deferral rate ie 19.4% [10]. In Singh P et al study (Uttar Pradesh) the deferral rate is

Table 5. Causes for temporary deferral

Causes for temporary deferral	2017	2018	2019	Total
Hypotension	15	14	16	45(15.05%)
Hypertension	20	18	13	51(17.06)
Low Hb	31	25	27	83(27.76%)
High Hb	2	4	5	11(3.68%)
Vaccination	15	11	9	35(11.71%)
Taken antibiotics	7	4	6	17(5.69%)
Dental extraction	4	2	3	9(3.01%)
Intake of alcohol	4	3	3	10(3.34%)
Major and minor surgery	2	2	2	6(2%)
Typhoid fever	1	1	1	3(1%)
Chicken pox	1	2	0	3(1%)
Tattooing	2	3	3	8(2.68%)
Asthma	1	0	1	2(0.67%)
Skin allergy	1	3	1	5(1.67%)
Previous donation not complete 3 months	1	1	1	3(1%)
Under weight below 45	1	1	0	2(0.67%)
Thin vein	1	0	1	2(0.67%)
Menstruation	1	0	0	1(0.33%)
Dengue	1	0	0	1(0.33%)
Ear piercing	1	0	0	1(0.33%)
Jaundice	0	0	1	1(0.33%)
	112	94	93	299

16.4% [11]. The deferral rate in Chauhan et al study (Uttarpradesh) is comparable to our study which is 5.56% [12]. In Harjot Kaur et al study (Punjab) the deferral rate is 6.99% [13]. A similar deferral rate is seen in Chenna et al study (Karnataka) 5.6% [14]. Vimal et al study (South India) show the deferral rate as 14.87% which is very much higher than our study [15]. In Nagarekha study (Karnataka) the deferral rate is only 4.27% [16]. In Unnikrishnan et al (Mangalore) the deferral rate is 5.2% [17]. The difference in deferral could be attributed due to difference in socioeconomic status, culture and the deferral criteria prevailing in various areas. In our study males were more common registered donors but the deferral rate is much higher in females. This is similar to most other studies [10,12,15]. In contrast to our study Nagarekha in her study showed males were more commonly deferred than females [16]. Anemia is more prevalent in females and this could be the cause. Females are needed to be more educated and motivated for blood donation. Permanent deferral (17.86%) is less common than temporary deferral (82.14%) in our study. This is comparable with many other studies [8,12]. Some studies show a low rate of temporary deferral [13]. Henshaw et al in their study revealed permanent deferral is more common [9]. Among permanent deferral high risk behavior particularly seropositive for hepatitis B is the most common cause (52.30%). The seropositive

rate in our study is 0.82%. Out of 6701 cases only 55 were serology positive. All the serology positive cases were males and all were replacement donors in our study. This emphasises the need for recruitment of voluntary donors. It is comparable to Unnikrishnan et al study (1.58%) This is followed by HCV and medications. This is in line with other studies [9,10,13,15]. But in Nagarekha and Chauhan et al study accounted hypertension as the most common cause of permanent deferral [16]. In Tufail et al study Hepatitis C positivity is the most common cause for permanent deferral [8].

Among temporary deferral Anemia is the most common cause (27.75%) followed by hypertension. This is on par with other studies [8,9,10,12,13,15]. This reflects the socioeconomic status of this region and people's ignorance about their health. In our country prevalence of both clinical and subclinical anemia is high. This indicates the importance of implementation of awareness and screening programs of anemia so that the health of the donor is guarded and we can maintain a healthy and safe donor pool. Most of the donors who were deferred due to anemia had Hemoglobin concentration around 11 to 12 gm. By reducing the Hemoglobin cut-off value by 1gm we can pave a way to get some more donors and this will enable us to meet our country's rising demand for blood [18]. Repeated donors are at

risk of iron depletion hence it is essential to supplement them with iron to maintain their iron reserve [19]. Hypertension could be due to anxiety, fear of phlebotomy and white coat hypertension. These individuals should be handled with care and helped to become more comfortable and relaxed by describing the donation in a detail and simpler manner. Hypotension was the third most common cause of temporary deferral in our study since syncope is more commonly anticipated in them due to vasovagal attacks [20].

The other major causes of temporary deferral in decreasing order of frequency include vaccination, medication, intake of alcohol, tooth extraction and tattooing. These temporarily deferred individuals should be educated and motivated so that they return for donation after the period of deferral. The deferral rate among voluntary and replacement donors is 4.71% and 11.62% respectively. In accordance with NACO we don't entertain professional donors in our institute and recruit more voluntary donors by organizing camps [21]. Rehman and Jawaid from their study revealed voluntary blood donors are safer than replacement donors [22]. Unnikrishnan et al also showed a similar pattern [17] but Nagarekha in her study revealed voluntary donor deferral was more common than replacement donor deferral [16]. Henshaw et al study showed a higher rate of deferral for commercial and replacement donors [9]. By analyzing the donor deferral pattern we can understand that the knowledge about donor selection criteria is inadequate in common people. By educating them we can get a better acceptable rate and individuals with more tendencies to return for a future donation and less bitter feeling towards rejection.

5. CONCLUSION

The deferral rate in our study was 5.19%. Females should be educated and motivated for blood donation. As anemia is more common in them there should be more national programs to manage it more effectively. As temporary deferral was more common this necessities the education, motivation and follow up of temporary deferred donors so they could return for future donation.

CONSENT

As per international standard or university standard, Participants' written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

Institutional ethical clearance was obtained before the commencement of the study (745/IHEC/12-19).

ACKNOWLEDGEMENT

We acknowledge the management of CHRI, technicians and all the donors who are involved in the study

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Department of AIDS Control Ministry of Health and Family Welfare Government of India. Annual report 2008-2009. 2009; 27.
2. World Health Organization. Blood donor selection: guidelines on assessing donor suitability for blood donation. Geneva: World Health Organization. 2012;16–23.
3. GDBS Summary Report 2016: WHO global database on blood safety (GDBS) and blood safety indicators. Geneva: World Health Organization. 2016;2-17.
4. Gillet P, Neijens E. An original approach to evaluating the quality of blood donor selection: checking donor questionnaires and analyzing donor deferral rate. *Front Med.* 2018;5:74.
5. GDBS Summary Report 2009: WHO Global Database on Blood Safety (GDBS) and Blood Safety Indicators. Geneva: World Health Organization.
6. Siromani, Umakanth, Thasian T, Rita Isaac, Dolly Daniel, Selvaraj K G, Joy John Mammen, and Sukesh Chandra Nair. "Shortage of blood! youths, the trend setters, can they help to meet the needs?" *Natl J Med Res.* 2013;3(2):196-197.
7. Patel S, Patel J, Patel A, Pandya AN, Raja K, Dobariya G, et al. "The study of deferred blood donors at tertiary level hospital based blood bank of south Gujarat". *J Evolution Medic Dental Sci.* 2015;4(27):4590-98.
8. Samina Tufail, Fatima Babar, Nadeem Ikram, Masooma Raza, Huma Abdul-Shakoor Blood Donors Deferral- Causes Journal of Rawalpindi Medical College (JRMC). 2013;17(1):119-121

9. Henshaw Uchechi Okoroiwu and Enosakhare Aiyudubie Asemota Blood donors deferral prevalence and causes in a tertiary health care hospital, southern Nigeria BMC Health Services Research. 2019;19:51
10. Laila Al Shaer Ranjita Sharma Mahera AbdulRahman Analysis of blood donor pre-donation deferral in Dubai: characteristics and reasons Journal of Blood Medicine. 2017;8:55–60.
11. Singh P, Jaiswal VK, verma N et al. Pattern of blood donor deferral in a tertiary hospital of western U.P.: A five-year study review. Rec Adv Path Lab Med. 2016;2(3&4):16-19.
12. Chitra Chauhan, Rashmi Chauhan, Seema Awasthi, S. Dutta, Himanshu Joshi Pattern and outcome of donor deferral -? need of hour Int J Res Med Sci. 2018;6(1):289-292
13. Harjot Kaur, Rahul Mannan, Mridu Manjari, Sonam Sharma2, Tania Garg and Tejinder Bhasin Blood Donor Deferral Analysis: A Tertiary Care Centre Experience Annals of Pathology and Laboratory Medicine. 2016;03(05(Suppl)):471-75
14. Chenna D, Shastry S, Murugesan M, Baliga PB. Implication of deferral pattern on the donor pool: Study at a Tertiary Care Hospital. J Appl Hematol. 2015;6:111-14
15. Vimal M, Sowmya S, Nishanthi A, Ramya G. Evaluation of blood donor deferral causes: a retrospective study from South India. Annals Pathol Lab Medic. 2016;3(6):A605-611.
16. Nagarekha Kulkarni Analysis Of Donor Deferral In Blood Donors Journal of Evolution of Medical and Dental Sciences. 2012;1(6):1081-86
17. Unnikrishnan B, Rao P, Kumar N, Ganti S, Prasad R, Amarnath A, Reshmi B, Kaur V, Kesharwani P, Seetha M, Nautiyal A, Goel P, Aggarwal A. Profile of blood donors and reasons for deferral in coastal South India. Australasian Medical Journal. 2011;4(7):379-385
18. Gandhi MJ, Duffy K, Benike M, Jenkins S, Stubbs JR. Effect of increasing hemoglobin cutoff in male donors and increasing interdonation interval in whole blood donors at a hospital-based blood donor center. Transfusion (Paris). 2012;52(9): 1880–8.
19. Røsvik AS, Hervig T, Wentzel-Larsen T, Ulvik RJ. Iron status in Norwegian blood donors: comparison of iron status in new blood donors registered in 1993-1997 and in 2005-2006. Vox Sang. 2009;96(1):49–55.
20. Crocco A, D'Elia D. Adverse reactions during voluntary donation of blood and/or blood components. A statistical-epidemiological study. Blood Transfus. 2007;5(3):143–52
21. NACO. An action plan for blood safety, MOHFW, GOI, [Online];2007. Available: <http://www.nacoonline.org/upload/Final%20Publications/Blood%20Safety/An%20Action%20Plan%20for%20blood%20safety.pdf>.
22. Rehman M and Jawaid SA. Need for a national blood policy to ensure safe blood transfusion. Pak J Med Sci. 2004;20(2):81-84.

© 2021 Thivya et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/76668>