



Enhancing Timber and Medicinal Plant Resources: Propagation Techniques for *Khaya senegalensis* (Desr.) A. Juss. in Tropical Regions

**Yao Kouakou Abessika Georges ^{a*},
Gbotto Ahou Anique ^a, Gnamien Yah Gwladys ^b,
Kouame Brou Grâce Emmanuella ^a,
Akaffou Doffou Sélastique ^a and Zoro Bi Irié Arsene ^c**

^a Department of Agronomy and Forestry, Laboratory of Genetics, University of Jean Lorougnon Guede, BP 150 Daloa, Côte d'Ivoire.

^b Department of Agronomy and Forestry, Laboratory of Plant Physiology, University of Jean Lorougnon Guede, BP 150 Daloa, Côte d'Ivoire.

^c Laboratory of Biology and Improving Vegetable Productions, Natural Sciences Unit, University Nangui Abrogoua (UNA), BP 801 Abidjan, Côte d'Ivoire.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jeai/2024/v46i72584>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/118147>

Short Research Article

Received: 02/04/2024

Accepted: 05/06/2024

Published: 13/06/2024

*Corresponding author: E-mail: georgesyao@yahoo.fr;

ABSTRACT

Human uses the resources of tropical forests, which contain great biological diversity, for food, traditional and commercial purposes, including on an international scale. Some species have been exploited for timber production. The aim of this work was to identify the best part of the shoot for regeneration of *Khaya senegalensis* in the two seedling substrates. This work will provide a database for the regeneration and domestication of forest products, with *Khaya senegalensis* as a model plant. It is one of the medicinal and timber species on which sustained attention and priority actions should be focused. The main objective of this study was to determine whether this plant, with its high health value, can be regenerated by cuttings. The set-up comprised 180 bags arranged in a random block of 90 bags for each of the seeding substrates (black earth and red earth). The results showed that regeneration by cuttings is possible in the substrates. Old cuttings showed the best viability rates, at 70% and 43.33% for black earth and red earth substrates respectively. The best regeneration percentage was observed for aged cuttings (36.66% and 20%). These results showed variability in length and collar diameter according to cutting type and substrate. Of the three types of explants tested, the old part of the stem proved better than the young and intermediate parts. However, it would be interesting to improve the rate of regeneration and rooting of cuttings by using vegetative hormones.

Keywords: *Khaya sénégalensis*; domestication; cuttings; forest biodiversity; regeneration.

1. INTRODUCTION

Humans have always used the resources of tropical forests, which contain a wealth of biological diversity (Sosef et al. [1]), for food, traditional and commercial purposes, including on an international scale (De Wasseige et al.[2]). Some species have been exploited for timber production (Sawyer [3]). *Khaya senegalensis* falls into this category. However, overexploitation of the species has led to its inclusion on the International Union for Conservation of Nature's red list (Adjahossou et al.[4]) as a species facing a very high risk of extinction in the wild (IUCN [5]). This species produces one of the most prized woods on African continent. It has been marketed under the name Mahogany for around two centuries. *Khaya senegalensis* is one of the medicinal and timber species that require sustained attention and priority action on account of its socio-economic importance (Kantende et al. [6]). It is a multi-purpose species and one of the largest and most majestic trees in the Sudanian region (Eyog [7]). Various organs of the species are used for their medicinal properties (Kolawole et al.[8]). Indeed, *K. senegalensis* is a tropical evergreen tree whose organs are widely used in the treatment of several diseases (Sokpon & Ouinsavi [9], Nikiema & Pasternak [10]; Adomou et al.[11]). The leaf, bark, fruit and root have highly recognized medicinal, food and ornamental values. Oil from the seeds is used in African cooking. Leaves and bark are used to treat stomach ache, diarrhea and constipation. These

organs also treat malaria. They are a remedy for jaundice and leprosy [12,13]. It eliminates intestinal parasites, and is a plant with febrifuge, antibiotic and vermifuge therapeutic properties. In addition to its medicinal importance, its wood is also heavily exploited as timber and service wood, which poses threats to the survival of this species, as it faces abusive exploitation by the rural population (Dainou et al. [14]). This results in forest degradation and loss of biodiversity. However, to remedy this situation, the domestication of this medicinal species with high therapeutic or nutritional value is worth considering. The general objective of this work is to regenerate *K. senegalensis* by cuttings. Specifically, the aim is to determine the best part of the stem to use as a cutting for good regeneration, and to determine the seeding substrate that conditions the best regeneration of the cuttings.

2. MATERIALS AND METHODS

2.1 Experimental Site

The study took place in one of the experimental nurseries of the University Jean Lorougnon GUEDE located in the town of Daloa at 6°53 north latitude and 6°27 west longitude. The town is located in the Haut Sassandra Region in West-Central Côte d'Ivoire. It is around 141 km from Yamoussokro, the political capital, and around 400 Km from Abidjan, the economic capital (Diomandé et al. [15]). Daloa is bordered to the north by the town of Vavoua, to the south by Issia, to the west by Duekoue and to the east by

Bouafle. The vegetation, which belongs to the mesophilic sector, is largely made up of dense forest, which has now disappeared to make way for various cash crops (Sangare et al.[16]). The soil is ferrallitic and the climate is humid tropical, with one rainy season and two dry seasons (N'guessan et al.[17]). Dry and wet seasons alternate with temperatures ranging from 24.65°C to 27.75°C on average.

2.2 Plant Material

The plant material used consists of *Khaya senegalensis* cuttings. These cuttings consist of three parts of the stem. The old part, which corresponds to the old wood directly connected to the root; the growing part, characterized by the part that generally bears the leaves; and the middle part, which corresponds to the intermediate zone between the old wood and the young part.

2.3 Methods

2.3.1 Taking and preparing cuttings

Cuttings were taken from the stems of plants showing variability in stem length and diameter. Each cutting has at least one node and is 14 cm long, 10 cm below the node and 4 cm above. Each stem was subdivided into three parts (old, young and middle). A total of 30 cuttings of each type were taken.

2.3.2 Seeding substrate

Two substrates commonly used for cuttings were tested. These were red soil taken from the Jean

Lorougnon GUEDE University in Daloa, and black soil taken from a former garbage dump.

2.3.3 Experimental design

The trial was conducted in a greenhouse built from the branches of oil palms (*Elaeis guineensis*), with wires used to connect the branches and a transparent tarpaulin to cover the nursery. The dimensions of the greenhouse are 9.54 m long, 3.3 m wide and 2.9 m high. The set-up comprised 180 bags arranged in a block of 90 bags for each of the seeding substrates. The cuttings were planted at a depth of 5 cm in the substrate. Each substrate received a total of 90 cuttings, i.e. 30 cuttings of each type, including 30 old cuttings, 30 middle cuttings and 30 young cuttings. The cuttings are watered every two days and hand weeding is carried out as soon as weeds appear in shaded areas.

2.3.4 Observed parameters

The parameters observed are bud break rate and time, bud height measured every week, bud neck diameter, regeneration and cuttings mortality rate. For this purpose, cuttings having developed a new leafy stem are counted at the end of the experiment, while the mortality rate concerns the proportion of dead cuttings no longer capable of developing buds. At the end of the experiment, the total number of each type of dead cuttings was determined according to the substrate. Observations lasted two months from April to June 2021.




Fig. 1. Type of cuttings a .Older part ; b .Middle part ; c .Younger part

2.3.5 Statistical analysis of data

The various parameters collected were subjected to statistical analysis. Two factors (substrates and cuttings) were subjected to an analysis of variance. When a difference was observed for a parameter tested for a given factor. The ANOVA was completed by the small significant difference test (SSDT). This test highlights the best parameter(s) tested. The data collected were analyzed using STATISTICA version 7.1 software.

3. RESULTS AND DISCUSSION

3.1 Results

3.1.1 Bud development and regeneration of cuttings

Cutting regeneration begins with bud formation. The bud formed develops over time into a leafy stem. In the greenhouse, we were able to follow this process. The bud forms on the 7th day after cultivation in the various substrates, and the first

leaves of the young stem appear on the 13th day. The leafy stem evolves over time.

Table 1 shows the results of the multivariate analysis of two factors studied: substrate ($P=0.046$) and cuttings used ($P<0.001$). This means that the results obtained are significantly influenced by the nature of the substrate and the type of cuttings used. Thus, in this work, results are presented according to substrate and type of cutting.

3.1.2 Mortality and viability of cuttings on different types of substrate

After 40 days' observation, T1 cuttings showed the best viability rates, on substrates. T2 cuttings showed low viability rates on the substrates. T3 cuttings recorded 100% mortality, the lowest viability rate in both substrates (Table 2). No young cuttings survived on either substrate. The red earth substrate thus recorded the highest mortality rate, while the black earth substrate recorded the highest viability rate (Table 2).




Fig. 2. Bud development, a : Formation of axillary buds, b : Development of petioles and appearance of the first leaves on the stem, c : Formation of leaves around the young stem

Table 1. Multivariate analysis of significance using Wilk's test

| | Test | Value | F | dl effect | P |
|---------------------|------|-------|--------|-----------|-------|
| ord.origin | Wilk | 0.872 | 12.637 | 2 | 0.000 |
| Substrates | Wilk | 0.965 | 3.136 | 2 | 0.046 |
| Cuttings | Wilk | 0.844 | 7.594 | 4 | 0.000 |
| Substrates*Cuttings | Wilk | 0.940 | 2.701 | 4 | 0.031 |

Table 2. Viability and mortality rates of cuttings according to substrate type

| Substrate | Red earth | | | Black earth | | |
|-----------------------|-----------|----|-----|-------------|-------|-----|
| | T1 | T2 | T3 | T1 | T2 | T3 |
| Type of cuttings | | | | | | |
| Viability (%) | 43.33 | 10 | 0 | 70 | 16.66 | 0 |
| Mortality (%) | 56.66 | 90 | 100 | 30 | 27 | 100 |
| Overall viability (%) | 17.7 | | | 28.887 | | |

T1 Older part ; T2 Middle part ; T3 younger part

Table 3. Regeneration rate of cuttings according to substrate

| Substrates | Regeneration of cuttings | | |
|-------------|--------------------------|-------|-------|
| | T1(%) | T2(%) | T3(%) |
| Red earth | 20 | 6.67 | 0 |
| Black earth | 36.67 | 10 | 0 |

T1 : Older part ; T2 : Middle part ; T3 : younger part

Table 4. Average stem length and average collar diameter depending of substrate

| Cuttings | Stem length (cm) | | | Collar diameter (cm) | | |
|-------------|----------------------------|---------------------------|----------------------|---------------------------|--------------------------|----------------------|
| | T1 | T2 | T3 | T1 | T2 | T3 |
| Red Earth | 19.667 ^a ±1,461 | 9.00 ^b ±2.53 | 0 ^c ±0.00 | 0.45 ^a ±0.063 | 0.40 ^a ±0.11 | 0 ^b ±0.00 |
| Black Earth | 12.272 ^a ±1,079 | 10.333 ^a ±2.06 | 0 ^b ±0.00 | 0.463 ^a ±0.046 | 0.40 ^a ±0.089 | 0 ^b ±0.00 |
| F | 33.047 | 6 | | 0.334 | | |
| P | P< 0.001 | | | P< 0.001 | | |

T1 : Older part ; T2 : Middle part ; T3 : younger part

3.1.3 Regeneration of cuttings in different types of substrate

Regeneration rates varied in the substrates. For different substrates, red earth and black earth, the highest percentage of regeneration was observed in T1 cuttings, followed by T2 cuttings (Table 3). As for T3 cuttings, we observed a total absence of regeneration. The regeneration rates observed in Table 3 show that black earth is a better substrate for regenerating cuttings.

3.1.4 Variation in mean stem length and collar diameter

The length and collar diameter of regrowths varied according to the type of cuttings and the nature of the substrate. No observations were made on young cuttings which did not regenerate. The results of the statistical analyses presented in Table 4 showed a significant difference in the length and collar diameter of regrowths on the cuttings. However, the nature of the substrate did not show any significant differences in the length and collar diameter of the regenerated cuttings.

3.2 Discussion

The results of this work showed that young cuttings had a low regeneration rate. In fact, young stems have excellent cell division qualities, but once cut, they contain fewer nutrient reserves. These portions of stem separated from the mother stem, no longer receiving reserves from elsewhere, dry out, leading to the death of the cuttings. Our results are in line with the work of Traoré [18], who reported that the inability of young *Piliostigma reticulatum* cuttings to regenerate was due to the fact that they did not contain sufficient nutrient

reserves to allow eventual emergence. On the other hand, according to Hartmann et al. [19], these young cuttings were subjected to desiccation and death. In addition, the same authors revealed that higher ambient temperatures could lead to bud break before root initiation, and increase water loss through transpiration. This will very often lead to cuttings drying out and dying [20,21].

As far as cuttings are concerned, the older part of the *Khaya senegalensis* stem was the best part for regenerating cuttings (Auneerudy and Pinglo [22]). Taking cuttings from the aged part of the *K. senegalensis* stem gave satisfactory results, since by the sixtieth day of the trial, some cuttings had produced root buds and others true roots for shoot survival and development. Regeneration of old cuttings therefore took place over a long period. Indeed, the regeneration capacity of aged cuttings could be explained by their ability to retain the nutrients required for bud formation and development over a long period. However, some cuttings can also develop leafy stems, but without a root system. Similar results were obtained with *Guiera senegalensis* cuttings in Burkina Faso (Bationo [23]). This author reports that cuttings without root systems are kept alive by accumulated reserves, but gradually dry out as these reserves are depleted.

This study also demonstrated the influence of substrate in the regeneration of *K. senegalensis* cuttings. The highest mortality and lowest viability of cuttings were observed in the red soil substrate, while the black soil induced the lowest mortality and highest viability. On the one hand, this result could be explained by the fact that red earth is a very clayey soil, with the capacity to retain water (Soro [24]). This capacity leads to an

excess of water, and the massive humidity in the substrate will result in rotting of the cuttings. This will lead to the death of the cuttings. On the other hand, red soil compacts easily, according to Soro [24], preventing the roots from breathing. This process will lead to asphyxiation of the roots, which could also lead to the death of the cuttings.

With regard to growth parameters, the results showed that the nature of the substrate had no statistical influence on the evolution of growth parameters. These results corroborate those obtained in the work of Silué et al.[25] who showed that there was no significant difference in the growth of *Khaya senegalensis* seedlings tested in different soil types. However, analysis of the effect of different substrates on the regeneration qualities of cuttings revealed that the black earth substrate induced the best results for growth parameters in *K. senegalensis*. These results are similar to those of Silué et al.[25] and Onana [26], who observed better values for growth parameters in *Khaya senegalensis* and *Pericopsis elata* respectively on typical substrates. The experimental period and conditions under which the trials were carried out may also explain the overall results obtained.

4. CONCLUSION

The main objective of this study was to determine the potential of *K. sénégaleensis*, a plant with a high health value, to be regenerated by cuttings. Following experiments with different types of cuttings (old, mature and young) on different substrates (black earth and red earth), the results showed that the type of substrate did not influence the regeneration rate, but rather the mortality rate. Of the three types of explants tested, the aged part or old part proved to be the better part of the stem than the young and intermediate parts. Although there was no significant difference between the substrate and the growth parameters, the black soil showed the best conditions for viability and regeneration by cuttings of *K. senegalensis*. It would therefore be interesting to improve the regeneration rate and rooting of cuttings through the use of vegetative hormones. In addition, a genotype effect test could be carried out to propose in-vitro culture for recalcitrant genotypes.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image

generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Sosef M, Dauby G, Blach-Overgaard A, van der Burgt X, Cata-rino L, Damen T. Exploring the floristic diversity of tropical Africa. BMC Biology. 2017;15:15. Available:<https://doi.org/10.1186/s12915-017-0356-8>
2. De Wasseige C, de Marcken P, Bayol N, Hiol H F, Mayaux Ph, Desclée B. The forests of the Congo basin. State of the Forests 2010. Luxembourg, Publications Office of the European Union. 2012;276. Available:https://www.observatoire.comifac.net/docs/edf2010/FR/Etat-des-forets_2010.pdf
3. Sawyer J. The Timber Trade. In: Sayer JA, Harcourt CS, Collins NM. (eds). The atlas for the conservation of African tropical forests. French edition, Doumenge C. (ed.). Paris, France, Jean-Pierre de Monza and IUCN France.1996;55-60
4. Adjahossou SGC, Yaoitcha A, Houehanou D. T, Gouwakinnou G N, Sode AI, Houinato M, Sinsin B. Favorable habitats for the conservation of *Khaya senegalensis* in Benin. Data sheets. National Library, Benin. 2018;12.
5. IUCN. IUCN Red List categories and criteria. <http://www.iucnredlist.org/> (accessed 05/15/2017) 2017.
6. Kantende AB, Birnie A, Tengas B. Useful trees and shrubs for Uganda. Regional Soil Conservation Unit RSCU/SIDA, Nairobi, Kenya ; 1995
7. Eyog M O. Monographic study of Caïlcédra; *Khaya senegalensis* (Desr) A. Juss. (Meliaceae). Nkolbisson Forestry Research Center, Maroua Branch, Cameroon. 1987;26.
8. Kolawole OT, Kolawole SO, Ayankunle AA, Olaniran OI. - Anti-hyperglycemic effect of *Khaya senegalensis* Stem bark aqueous extract in Wistar Rats. European Journal Medicinal Plants. 2012;2(1):66-73.
9. Sokpon N, Ouinsavi C. Management of *Khaya senegalensis* plantations in Benin. Woods and Forests of the Tropics. 2004;279:37-46.

10. Nikiema A, Pasternak D. *Khaya senegalensis* (Desr.) A.Juss. PROTA (Plant Resources of Tropical Africa).
11. Adomou A, Yedomonhan H, Djossa B, Legba S, Oumorou M, Akoegninou A. Ethnobotanical study of medicinal plants sold in the Abomey-Calavi market in Benin. *International Journal of Biological and Chemical Sciences*. 2012;6:745-772
12. Ocholla JA, Inoti SK, Obwoyere GO. Comparative Assessment of Regeneration, Structure and Species Diversity of Woody Vegetation in Disturbed and Undisturbed Sites of a Secondary Montane Forest, Kenya. *Asian Journal of Research in Agriculture and Forestry*. 2023;9(1):38–48. <https://doi.org/10.9734/ajraf/2023/v9i1195>
13. Wagner RG, Gonzalez-Benecke CA, Nelson AS, Jacobs DF. Forest regeneration in changing environments. *New Forests*. 2018, Nov;49: 699-703.
14. Daïnou K, Blanc-Jolivet, Degen B, Kimani P, Ndiade-Bouroubou D, Donkpegan ASL. Revealing hidden species diversity in closely related species using nuclear SNPs, SSRs and DNA sequences – a case study in the tree genus *Milicia*. *BMC Evolutionary Biology*. 2016;259. Available:<https://doi.org/10.1186/s12862-016-0831-9>
15. Diomandé M, Kouamé KB & Koko AC. Comparison of the chemical properties of peanut and cashew oil and meal on the Daloa market, Ivory Coast. *Engineering and Applied Sciences*. 2017;4(11):28-32.
16. Sangare A, Koffi E, Akamou F, Fall CA. State of plant genetic resources for food and agriculture. Second national report. Ministry of Agriculture, Republic of Côte d'Ivoire. 2009;64.
17. N'guessan C A, Abo K, Fondio L, Chiroleu F, Lebeau A, Poussier S, Wicker E, Koné D. So Near and Yet so Far: The Specific Case of *Ralstonia solanacearum* Populations from Côte d'Ivoire in Africa . *Bacteriology*. 2012;102:733-740.
18. Traore M. Study of the phenology, natural regeneration, and uses of *Piliostigma reticulatum* Hochst. In the North Sudanese zone of Burkina Faso: Rural Development Engineer's dissertation; Polytechnic University of Bobo Dioulasso. 2000;95.
19. Hartmann HTT, Kester DE, Davies FT, Geneve Jr RL. *Plant propagation principles and practices* 6th edition, pritice-hall in somon et Schuster, upper saddle River, New Jersey 07458, USA. 1977;770.
20. Garbaye J, Tacon F. Production of oak and beech plants from herbaceous cuttings. French Academy of Agriculture, extract from the minutes of the meeting of June 21. 1978;7.
21. Mapongmetsem MP, Djoumessi M, Fawa G, Doumara DG, Noubissie TJ, Tientcheu A. Domestication of *Vitex doniana* Sweet (verbenaceae): influence of substrate type, hormonal stimulation, leaf surface and node position on the rooting of uninodal cuttings 23 p. *Journal of Agriculture and Environmental for International Development*. 2012;106:23-45.
22. Aumeerudy Y, Pingolo F. Phytopractice in tropical regions. Aprimary survey of traditional crops improvement techniques. UNESCO/MAB/Tropical Botanical Laboratory. Montpellier. 1989;71.
23. Bationo BA. Study of agroforestry potential, multiplication and uses of *Guiera senegalensis*, Engineering dissertation, Institute of Rural Development, University of Ouagadougou. 1994;67.
24. Soro B. Soil classification of Côte d'Ivoire, 2018.
25. Silue, PA, Koffi KAD, Koffi, AB, Kouassi, KE. Germination tests and monitoring of growth performance of *Khaya senegalensis* (Desv.)A plants. Juss., in the Sudanian zone (Ivory Coast). *Journal of Animal and Plant Sciences*. 2021; 48(2):8673-8685
26. Onana MH. Silviculture and natural regeneration of *Pericopsis elata* (Harms) Var. Meeuwen in plantation: Case of the Eastern and Southern regions. Master's degree in Water, Forests and Hunting, University of Dschang, Cameroon, 2009; 88.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/118147>