



# Effect of Different Fertilizer and Silicon Levels on Grain Quality of Maize (*Zea mays* L.)

Seema Pooniyan <sup>a</sup>, K.K. Yadav <sup>a</sup>,  
Tushar Kumar <sup>a\*</sup> and Neha Meena <sup>b</sup>

<sup>a</sup> Department of Soil Science and Agricultural Chemistry, Rajasthan College of Agriculture (MPUAT), Udaipur, Rajasthan, India.

<sup>b</sup> Department of Soil Science and Agricultural Chemistry, Agriculture University, Kota, Rajasthan, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ijpss/2024/v36i84931>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/121417>

Original Research Article

Received: 10/06/2024

Accepted: 12/08/2024

Published: 19/08/2024

## ABSTRACT

A field experiment with split plot design was conducted to study the 'effect of fertility and silicon levels on the nutritional quality of maize' during *Kharif* 2022 and 2023 at Rajasthan College of Agriculture (RCA), MPUAT, Udaipur, Rajasthan, India. The main plot treatments consist of four levels of Compound NPK fertilizer (F<sub>0</sub>: Control, F<sub>50</sub>: 50% RDF, F<sub>75</sub>: 75% RDF and F<sub>100</sub>: 100 % RDF) and the sub-plot treatments consist of four levels of soil application of Diatomaceous Earth (Si<sub>0</sub>, Si<sub>100</sub>, Si<sub>200</sub> and Si<sub>300</sub>). Pratap Makka-9 of maize variety was used for research purpose. The protein content, protein yield, carbohydrate and energy in maize grain were higher under 100%

\*Corresponding author: E-mail: [tusharkumar557@gmail.com](mailto:tusharkumar557@gmail.com);

**Cite as:** Pooniyan, Seema, K.K. Yadav, Tushar Kumar, and Neha Meena. 2024. "Effect of Different Fertilizer and Silicon Levels on Grain Quality of Maize (*Zea Mays* L.)". *International Journal of Plant & Soil Science* 36 (8):992-998. <https://doi.org/10.9734/ijpss/2024/v36i84931>.

RDF than other levels of fertilizer. Application of 300 kg Diatomaceous Earth per hectare recorded higher protein content, protein yield, carbohydrate and energy in maize grain over the rest of silicon levels. The crude fat in maize grain was lower in 100% RDF. Application of 300 kg Diatomaceous Earth per hectare recorded lower crude fat in maize grain. Significantly higher protein yield was recorded under the combined application of 100% RDF and 300 kg diatomaceous earth per hectare. However, protein yield was at par in the application of 75% RDF with 300 kg Diatomaceous Earth per hectare and 100% RDF with 200 kg Diatomaceous Earth per hectare. The lowest crude fat was recorded under the combined application of 100% RDF and 300 kg Diatomaceous Earth per hectare over the rest of treatment combinations. Application of 100% RDF with 300 kg DE ha<sup>-1</sup> (F<sub>100</sub>Si<sub>300</sub>) was recommended treatment for nutritional high quality maize grain.

**Keywords:** maize grain; protein content and yield; crude fat; carbohydrate; diatomaceous earth.

## 1. INTRODUCTION

Maize (*Zea mays* L.) is the most efficient, productive cereal crop that has numerous industrial importance and ranks first in global production [1]. Maize is a staple food in many parts of the world, with total maize production exceeding that of wheat or rice production [2]. It is a good source of carbohydrates, protein, iron, vitamin B, and minerals. Approximately 70% of maize is used for animal feed, with the remainder going into industrial products [3]. Maize is an exhausting crop, thus the nutrient requirements cannot be met only by soil nutrient reserves; additional nutrients can be addressed through fertilizer application. Maize yields in Rajasthan are low because of an uneven nutrient application. Scientists face a challenge when recommending fertilizer doses since they must meet both the crop's nutritional need and the production system's sustainability [4].

Silicon is the most plentiful element (in the earth's crust) after oxygen, with soils containing roughly 28% silicon by weight [5]. Because of its abundance in the biosphere, silicon's importance as a micronutrient for higher plants is difficult to demonstrate. Agricultural activities tend to extract a substantial amount of silicon from soil. Even highly filtered water contains approximately 20 ppb Si Werner and Roth, [6], while the leaves of silicon accumulator plants which weren't exposed to silicon treatment typically contain between 0.5 and 1.9 mg Si g<sup>-1</sup> leaf dry weight. Although silicon is considered a necessary component for many Poaceae species, direct evidence that it is a component of a molecule that is a necessary plant constituent or metabolite is still lacking, making it impossible to prove that silicon is necessary for all higher plants [7].

## 2. MATERIALS AND METHODS

### 2.1 Location

The experiment was conducted during *kharif* season of 2022 and 2023 at Instructional Farm, RCA, MPUAT (Udaipur). The geographical coordinates of the experimental site is 24°35' N latitude and 73°42' E longitude at an altitude of 582.2 m above mean sea level. RCA, MPUAT, Udaipur comes under the IV a Zone of Rajasthan. The soil of the experimental site was clay loam soil (Typic Haplustepts).

### 2.2 Experimental Details

The treatments were laid out in a split plot design with four levels of fertilizer as control, 50% RDF 75% RDF and 100% RDF (F<sub>0</sub>, F<sub>50</sub>, F<sub>75</sub> and F<sub>100</sub>, respectively) with four levels of silicon @ control, 100, 200 and 300 kg DE ha<sup>-1</sup> (Si<sub>0</sub>, Si<sub>100</sub>, Si<sub>200</sub> and Si<sub>300</sub>, respectively). Levels of fertilizer were included as a main plot and silicon levels as sub plot. Recommended dose of fertilizer (RDF) @ 90: 60: 40 N: P<sub>2</sub>O<sub>5</sub>: K<sub>2</sub>O kg ha<sup>-1</sup> was used and were calculated as per treatments based on gross plot size. The variety of maize Pratap Makka-9 was used for sowing. Sowing were under taken on 15<sup>th</sup> July and 8<sup>th</sup> July during 2022 and 2023, respectively. The nutrients viz., nitrogen, phosphorous and potassium were applied in the form of urea, di-ammonium phosphate (DAP) and muriate of potash (MOP), respectively. The silicon levels were applied in the form of Diatomaceous Earth (DE).

DE is a sedimentary rock formed silica hydroxide minerals and is amorphous in nature, being porous and earthy chalk like mineral. This DE is primarily consists of the remnants of silica skeletons of diatoms. The distinctive chemical

composition of DE usually comprises of 86% as silicon 5% as Na, 3% as Mg and 2% as Fe, respectively. Diatoms are accrued in the sediment of rivers, streams, lakes, and oceans over a period of time and it is white gray, or off-white in color, and can contain water in the 3 to 13% range [8].

## 2.3 Analysis of Nutritional Quality of Maize

### 2.3.1 Protein content

The protein content was calculated by measuring the nitrogen content of the sample using the Kjeldahl method [9]. The protein content of the products was calculated by multiplying the nitrogen content of the material by a factor of 6.25 [10].

$$\text{Protein content (\%)} = \text{Percent Nitrogen} \times 6.25$$

### 2.3.2 Protein yield

Protein yield was computed by using following formula;

$$\text{Protein yield (kg ha}^{-1}\text{)} = \frac{\text{Seedyield (kg ha}^{-1}\text{)} \times \text{protein content (\%)}}{100}$$

### 2.3.3 Crude fat

The fat content of the maize samples was estimated as crude ether extract of moisture free samples using the Soxhlet's Extraction Method on Socs Plus System. A thimble was filled with a weighed amount of moisture- free sample (5 g) and placed in the thimble holder, which was placed in an already weighed beaker, and 80 ml petroleum ether (60-80 °C) was poured into the beaker. The system was loaded with beakers and the temperature was set to 100 °C. After 120 minutes of operation, which was twice the initial boiling temperature. Rinsing was thus performed twice to collect any remaining fat in the sample. Beakers were removed and placed in a hot air oven. The beakers were weighed after the thimble holders were removed. The fat content of the sample was determined using the following formula:

$$\text{Crude fat (\%)} = \frac{\text{Weight of ether extract fat (B-A)}}{\text{Weight of sample (g)}} \times 100$$

Where,

A= Weight of empty flask (g)

B= Weight of flask + fat (g)

B-A= Weight of fat (g)

### 2.3.4 Carbohydrate content

The phenol sulphuric acid method as described by Roberts et al., 2011 was used to estimate carbohydrate contents in selected maize accessions. The powder of each accession (0.1g) was mixed in 5 mL of 2.5 N HCL and heated in a water bath for 3 to 4 hours. It was then neutralized by adding sodium carbonate (Na<sub>2</sub>CO<sub>3</sub>) and volume increase up to 100 to 150 mL.

Glucose with good strength was used as a standard for carbohydrate estimation. Each experimental sample was mixed with 1 mL of 5% phenol solution and 5 mL of 96% sulphuric acid (H<sub>2</sub>SO<sub>4</sub>) solution and kept at 30 °C for 20 min. After that absorbance was taken at 490 nm of wavelength of spectrophotometer obtained from standard curve was used to estimate carbohydrate content in selected maize cob grain accessions.

$$X \frac{Y - 0.2981}{0.0337} \times 100$$

### 2.3.5 Energy

The sample's energy value were calculated using physiological energy values of 4 Kcal per gram for protein, 9 Kcal per gram for fat and 4 Kcal per gram for carbohydrate.

$$\text{Energy (Kcal/100g)} = [(\% \text{ protein} \times 4) + (\% \text{ carbohydrates} \times 4) + (\% \text{ fat} \times 9)]$$

## 3. RESULTS AND DISCUSSION

### 3.1 Effect of Fertility Levels

#### 3.1.1 Protein content

The data (Table 1) show it clear that rising fertility levels considerably improve the protein content of maize grain up to 75% RDF (F<sub>75</sub>). When 75% RDF was applied, the protein content increased by 15.9 and 6.1% in the pooled mean, compared to the control (F<sub>0</sub>) and F<sub>50</sub> level of fertility. In terms of statistics, the protein content between 75% RDF and 100% RDF remained at par. The addition of NPK fertilizers is an important practices for increasing protein content in maize. Nitrogen is directly involved in protein synthesis, phosphorus supplies energy for metabolic processes, and potassium promotes enzyme performance and nutrient transport. Together,

these nutrients insure that maize plants develop healthily, efficiently use nutrients, and yield grains with higher protein content. A similar study was given by Ashoka *et al.* [11].

### 3.1.2 Protein yield

Protein yield of maize is presented in Table 1. The maximum value 528.3 kg ha<sup>-1</sup> for protein yield was observed in application of 100% RDF. The significant improvement in protein yield with the application of fertility levels seems to be due to enhancing production of protein content and higher grain yield.

### 3.1.3 Carbohydrate content

The data clearly shows that rising fertilizer levels improved the carbohydrate content in maize grain, however, the increase was only statistically significant up to 75% RDF (Table 2). The carbohydrate content in maize grain is statistically at par under 75 and 100% RDF. Increased fertilizer levels may improve carbohydrate content in maize grain, resulting in better growth, higher yields and superior nutritional value. A similar study was observed by Kour and Singh [12].

**Table 1. Effect of fertilizer and silicon on protein content and protein yield in maize grain**

Treatments	Protein Content (%)			Protein Yield (kg ha <sup>-1</sup> )		
	2022	2023	Pooled	2022	2023	Pooled
<b>Fertility levels</b>						
Control	9.13	9.32	9.23	310.2	327.6	318.9
50% RDF	10.06	10.10	10.08	437.8	452.9	445.3
75% RDF	10.63	10.75	10.69	521.7	536.3	529.0
100% RDF	10.73	10.91	10.82	551.9	564.7	558.3
<b>S.Em.±</b>	<b>0.11</b>	<b>0.14</b>	<b>0.09</b>	<b>7.55</b>	<b>9.04</b>	<b>5.89</b>
<b>C.D. (P=0.05)</b>	<b>0.37</b>	<b>0.49</b>	<b>0.27</b>	<b>26.13</b>	<b>31.27</b>	<b>18.14</b>
<b>Silicon levels</b>						
Control	9.56	9.70	9.63	377.8	389.2	383.5
100 kg DE ha <sup>-1</sup>	9.99	10.13	10.06	433.8	447.8	440.8
200 kg DE ha <sup>-1</sup>	10.48	10.57	10.53	489.5	502.2	495.9
300 kg DE ha <sup>-1</sup>	10.52	10.67	10.60	520.4	542.2	531.3
<b>S.Em.±</b>	<b>0.10</b>	<b>0.10</b>	<b>0.07</b>	<b>9.79</b>	<b>8.94</b>	<b>6.63</b>
<b>C.D. (P=0.05)</b>	<b>0.30</b>	<b>0.29</b>	<b>0.20</b>	<b>28.57</b>	<b>26.08</b>	<b>18.84</b>

**Table 2. Effect of fertilizer and silicon on carbohydrate content, crude fat and energy in maize grain**

Treatments	Carbohydrate Content (%)			Crude Fat (%)			Energy (kcal/100g)		
	2022	2023	Pooled	2022	2023	Pooled	2022	2023	Pooled
<b>Fertility levels</b>									
Control	72.20	72.58	72.39	1.18	1.20	1.19	335.92	338.37	337.15
50% RDF	74.32	74.76	74.54	1.11	1.15	1.13	347.54	349.80	348.67
75% RDF	76.98	77.38	77.18	1.01	1.05	1.03	359.49	361.99	360.74
100% RDF	77.72	78.71	78.22	0.92	0.99	0.95	362.10	367.38	364.74
<b>SEm±</b>	<b>1.03</b>	<b>1.07</b>	<b>0.74</b>	<b>0.02</b>	<b>0.01</b>	<b>0.01</b>	<b>3.95</b>	<b>4.31</b>	<b>2.92</b>
<b>CD(P=0.05)</b>	<b>3.58</b>	<b>3.69</b>	<b>2.29</b>	<b>0.05</b>	<b>0.05</b>	<b>0.03</b>	<b>13.67</b>	<b>14.92</b>	<b>9.01</b>
<b>Silicon levels</b>									
Control	72.23	72.59	72.41	1.14	1.18	1.16	337.45	339.82	338.64
100 kg DE ha <sup>-1</sup>	73.56	74.36	73.96	1.11	1.15	1.13	344.21	348.31	346.26
200 kg DE ha <sup>-1</sup>	76.59	77.59	77.09	1.02	1.06	1.04	357.45	362.20	359.83
300 kg DE ha <sup>-1</sup>	78.85	78.89	78.87	0.94	0.99	0.97	365.93	367.21	366.57
<b>SEm±</b>	<b>1.02</b>	<b>1.00</b>	<b>0.71</b>	<b>0.01</b>	<b>0.01</b>	<b>0.01</b>	<b>4.15</b>	<b>4.09</b>	<b>2.91</b>
<b>CD(P=0.05)</b>	<b>2.96</b>	<b>2.92</b>	<b>2.03</b>	<b>0.04</b>	<b>0.04</b>	<b>0.03</b>	<b>12.12</b>	<b>11.94</b>	<b>8.29</b>

### 3.1.4 Crude fat

The crude fat content of maize grain decreased dramatically as fertilizer levels increased (Table 2). Lower crude fat in maize grain was recorded with the application of 100% RDF. Fertilizers promote healthier plants, improve growth and metabolism, and result in decreased crude fat content. The same findings were also reported by Shinde *et al.*, [13].

### 3.1.5 Energy

The maximum energy (364.74 kcal/100gm) was obtained from maize grain under the application of 100% RDF which at par with the 75% RDF (Table 2). Fertiliser application to maize crops is critical for increasing grain energy content, which leads to higher yields, better quality, and improved plant health.

### 3.1.6 Effect of silicon levels

Silicon in general acts as in the strengthening of cell wall by provided structural integrity to plants, it can also act as a potential source of quality as it reduces the strain produced by abiotic factors such as flooding, drought and heat stress. Silicon is also known for its impact in the modification of biosphere in and around the rhizospheric zone this leads to higher plant nutrient uptake and better hormonal health of microbes and plants (Kandil *et al.* [14], Asadpour *et al.* [15], Khan *et al.* [16])

### 3.1.7 Protein content

The data in Table 1 show that increasing fertility levels significantly improves protein content in maize grain. The highest protein content recorded under the application of 300 kg DE ha<sup>-1</sup> (Si<sub>300</sub>). However, protein content remained statistically at par with 200 (Si<sub>200</sub>) and 300 (Si<sub>300</sub>) silicon levels. The 200 kg DE ha<sup>-1</sup> (Si<sub>200</sub>) reported an increase of 9.3 and 4.6 per cent in the pooled analysis over the control (Si<sub>0</sub>) and 100 kg DE ha<sup>-1</sup> (Si<sub>100</sub>), respectively. These results are in conformity with the findings of Ahmed *et al.* [17], Bakhat [18] and El-Temsah [19].

### 3.1.8 Protein yield

The data shown in Table 1 shows that increasing silicon levels significantly increases the protein yield in maize. The highest protein yield was recorded under 300 kg DE ha<sup>-1</sup> (Si) application. Silicon applications can help plants grow

healthier and produce more protein by reducing stress and boosting physiological functioning.

### 3.1.9 Carbohydrate content

The carbohydrate content increased significantly with the successive increase in the levels of silicon up to 200 kg DE ha<sup>-1</sup> (Table 2). The carbohydrate content remained statistically at par under 200 (Si<sub>200</sub>) and 300 (Si<sub>300</sub>) kg DE ha<sup>-1</sup>. The silicon application results in healthier, more resilient plants with higher carbohydrate output. Similar studies were reported by Iqbal *et al.* [20].

### 3.1.10 Crude fat

The data clearly shows in Table 2 that increasing the levels of silicon caused a significant decrease in crude fat in maize grain. The lowest crude fat content in maize grain was observed under the 300 kg DE ha<sup>-1</sup> application. The application of silicon makes plants healthier, and more resilient, optimizes their growth and resource utilization, resulted in reducing crude fat content. These results conform to the findings of Artyszak, [21].

### 3.1.11 Energy

The data (Table 2) shows that significantly enhanced energy in maize grain with increasing levels of silicon up to 200 kg DE ha<sup>-1</sup> (Si<sub>200</sub>) [16,22]. The energy in maize grain remained statistically at par under 200 (Si<sub>200</sub>) and 300 (Si<sub>300</sub>) kg DE ha<sup>-1</sup>. The use of silicon significantly increases maize growth, stress tolerance as well as improves the energy content in the grain [23].

## 3.2 Combined Effect of Fertilizer and Silicon

### 3.2.1 Protein yield

Application of 100% RDF with 300 kg DE ha<sup>-1</sup> (F<sub>100</sub>Si<sub>300</sub>) recorded highest protein yield which was significantly increased over the rest of the treatment combinations but found at par with the application of 75% RDF with 300 kg DE ha<sup>-1</sup> (F<sub>75</sub>Si<sub>300</sub>) and 100% RDF with 200 kg DE ha<sup>-1</sup> (F<sub>100</sub>Si<sub>200</sub>) (Table 3).

### 3.2.2 Crude fat

Application of 100% RDF with 300 kg DE ha<sup>-1</sup> (F<sub>100</sub>Si<sub>300</sub>) was recorded lowest crude fat which was significantly increased over the rest of the treatment combinations (Table 3) [13,9].

**Table 3. Combined effect of fertilizer and silicon on protein yield and crude fat in maize grain**

F x Si	Protein Yield				Crude Fat			
	Si <sub>0</sub>	Si <sub>100</sub>	Si <sub>200</sub>	Si <sub>300</sub>	Si <sub>0</sub>	Si <sub>100</sub>	Si <sub>200</sub>	Si <sub>300</sub>
F <sub>0</sub>	247.96	319.15	353.43	355.09	1.32	1.23	1.17	1.03
F <sub>50</sub>	401.12	426.84	474.78	478.52	1.17	1.15	1.14	1.08
F <sub>75</sub>	425.32	508.99	540.48	641.03	1.10	1.09	0.99	0.94
F <sub>100</sub>	459.54	508.29	614.80	650.55	1.06	1.07	0.87	0.81
<b>SEm±</b>	13.25				0.020			
<b>CD (P=0.05)</b>	37.69				0.057			

#### 4. CONCLUSION

The use of different levels of fertilizers and silicon can considerably improve the nutritional value of maize grain. Fertilizers give nutrients required for growth and development, whereas silicon promotes plant health and stress tolerance (source). These together measures improve grain energy content, protein content, carbohydrate content, protein yield and crude fat. Application of 100% RDF with 300 kg DE ha<sup>-1</sup> (F<sub>100</sub>Si<sub>300</sub>) was recommended treatment for nutritional high quality maize grain.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Qu Z, Luo N, Guo J, Xu J, Wang P, Meng Q. Enhancing sustainability in the new variety-based low emergy system for maize production by nitrogen optimization. *Renewable and Sustainable Energy Reviews*. 2024;199:114471.
2. Verma SK, Kumar S, Deepshikha MG, Yadav DN. Food Grain Trade Prospect of India: Markov Chain Analysis; 2024.
3. Kumar R, Bhardwaj A, Singh LP, Singh G. Environmental and economical assessment of maize cultivation in Northern India. *Process Integration and Optimization for Sustainability*. 2024;8(1): 165-179.
4. Shankar MA, Umesh MR. Site specific nutrient management (SSNM): An approach and methodology for achieving sustainable crop productivity in dryland Alfisols of Karnataka. *Technical Bull. University Agriculture Science, Bangalore (India)*; 2008.
5. Lindsay WL. Chemical equilibrium in soil. John Wiley & Sons, New York, NY; 1979.
6. Werner D, Roth R. Silica metabolism. *Inorganic Plant Nutrition*. 1983;682-694.
7. Epstein E. Silicon. *Annual Review Plant Biology*. 1999;50:641-644.
8. Gokavi N, Jayakumar M, Mote K, Surendran U. Diatomaceous earth as a source of silicon and its impact on soil physical and chemical properties, yield and quality, pests and disease incidence of Arabica Coffee cv. Chandragiri. *Silicon*. 2021;13:4583-4600.
9. Snell FD, Snell CP. *Colorimetric methods of analysis*. 3<sup>rd</sup> Ed. D. Van Nostrand Inc. New York. 1949;2.
10. Sarkar S, Mondal M, Ghosh P, Saha M, Chatterjee S. Quantification of total protein content from some traditionally used edible plant leaves: A comparative study. *Journal of Medicinal Plant Studies*. 2020;8(4):166-170.
11. Ashoka P, Anand SR, Smitha R. Effect of macro and micronutrients with organics on growth, quality, yield and economics of Baby corn (*Zea mays* L.) in Tungabhadra command area. *Crop Research (Hisar)*. 2009;37(1/3):15-18.
12. Kour J, Singh AP. Effect of crop geometry, integrated nutrient management on growth, production and nutrient uptake of sweet corn (*Zea mays* L. Saccharata). *Indian Research Journal of Genetics & Biotechnology*. 2020;12(1):30-37.
13. Shinde SA, Patange MJ, Dhage SJ. Influence of irrigation schedules and

- integrated nutrient management on growth, yield and quality of *rabi* maize (*Zea mays* L.). *International Journal of Current Microbiology and Applied Sciences*. 2014;3 (12):828-832.
14. Kandil EE, Lamloom SF, Gheith ESM, Javed T, Ghareeb RY, Abdelsalam NR, Hussain S. Biofortification of maize growth, productivity and quality using nano-silver, silicon and zinc particles with different irrigation intervals. *The Journal of Agricultural Science*. 2023;161(3):339-355.
  15. Asadpour S, Madani H, Mohammadi GN, Heravan IM, Abad HHS. Improving maize yield with advancing planting time and nano-silicon foliar spray alone or combined with zinc. *Silicon*. 2020;1-9.
  16. Khan WUD, Sharif F, Naeem MA, Farooq MA, Siddiq Z, Imran M. Chitosan polymerized silica composite as a potential silicon source: Modulation on antioxidant enzymes, ionic homeostasis, and grain quality in maize plants under Na<sup>+</sup> Stress. *Journal of Plant Growth Regulation*. 2023; 42(4):2374-2388.
  17. Ahmed M, Hassen FU, Qadeer U, Aslam MA. Silicon application and drought tolerance mechanism of sorghum. *African Journal of Agricultural Research*. 2011;6 (3):594-607.
  18. Bakhat HFSG. Role of silicon in plasmalemma H-ATPase hydrolytic and pumping activity in maize (*Zea mays* L.). Ph.D. thesis. Justus Liebig University, Giessen, Germany; 2012.
  19. El-Temsah ME. Response of rice yield, its components and quality to silicon and boron foliar application. *Middle East Journal of Agriculture Research*. 2017;6(4): 1259-1267.
  20. Iqbal N, Khan NA, Umar. Photosynthetic inhibition under salinity challenged environment: An insight into regulation of rubisco. *Functional Genomics, Physiological Processes and Environmental Issues*. 2011;167.
  21. Artyszak A. Effect of silicon fertilization on crop yield quantity and quality - A literature review in Europe. *Plants*. 2018;7(3):54.
  22. Rajoo R. Maize the Golden Grain of Himachal Pradesh. In Kalyani Press, Ludhiana, India; 1998.
  23. Roberts R, Elias R. Determination of carbohydrate using phenol sulphuric acid method. In: *Food Analysis* (4th Ed). Nielson, S. (ed.): Springer; 2011.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/121417>