



Biometrical Relationship between Body Weight and Body Measurements of Black Bengal Goat (BBG)

**Md. Ahsan Habib^{1*}, Ambia Akhtar², A. K. Fazlul Haque Bhuiyan³,
Md. Panir Choudhury¹ and Most Farhana Afroz¹**

¹Bangladesh Livestock Research Institute, Savar, Dhaka-1341, Bangladesh.

²Department of Livestock Services (DLS), Bangladesh.

³Bangladesh Agricultural University (BAU), Mymensingh-2202, Bangladesh.

Authors' contributions

This work was carried out in collaboration among all authors. Author MAH performed the statistical analysis and wrote the first draft of the manuscript. Author AA collected data and checked first draft of manuscript. Author AKFHB designed the study, wrote the protocol and checked the final draft of the manuscript. Authors MPC and MFA checked analysis and managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/CJAST/2019/v35i230172

Editor(s):

(1) Dr. Ogunlade, Clement Adesoji, Lecturer, Department of Agricultural Engineering, Faculty of Engineering, Adeleke University, Ede, Osun State, Nigeria.

Reviewers:

(1) Mahieu Maurice, INRA, France.

(2) Adeyinka . O. Akintunde, University of Ilorin, Nigeria.

(3) Prem Prakash Dubey, Guru Angad Dev Veterinary and Animal Sciences University, India.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/48413>

Received 25 January 2019

Accepted 15 April 2019

Published 02 May 2019

Original Research Article

ABSTRACT

Aims: To develop regression equations for estimation of live weight from the external morphological measurements.

Study Design: CRD with non-orthogonal hierarchy.

Place and Duration of Study: This study was carried out at 3 rural village communities of Bhaluka Upazila in Mymensingh district of Bangladesh from 2009 to 2013.

Methodology: All animals were ear-tagged and maintained under semi-intensive management system with scheduled vaccination and de-worming. Data were collected from a baseline survey along with 3 progressive generations produced from the community foundation stocks. Body length, chest girth, wither height, hip height and body weights were measured in a same day and recorded individually from birth up to 15 month age. Body weights were measured in kilogram by a hanging

*Corresponding author: E-mail: ahsan.rony@yahoo.com;

spring balance and other morphological parameters were measured in centimeter by a measuring tape. The data were analyzed by "SPSS 17.0" statistical program.

Results: A total of seven regression models were adopted and analysis of variance showed that all models were fitted significantly ($p < 0.001$). The correlation coefficient was higher when multiple body measurements were included in the model. The study also revealed that when chest girth as a single body measurement was included in the model gave the highest correlation coefficient ($R = 0.92$). For including multiple measurements, body length and chest girth are the best external body measures which exposed same correlation coefficient ($R^2 = 0.92$) when included more than these two body measures in the model. The differences between actual body weight and body weight predicted from regression equation for different ages were less than 1% ($p > 0.05$) and correlation coefficient between weights was 0.92 ($p < 0.01$).

Conclusion: There are strong correlations among morphometric body measurements. Thus, body weight can be estimated from a single or multiple body measurements by regression equation. Chest girth is the best single predictor for estimating live body weight with high accuracy.

Keywords: Black Bengal goat; morphological measurements; biometrical relationship; regression equations.

1. INTRODUCTION

In Bangladesh there is only one native goat breed commonly known as the Black Bengal Goat (BBG) and found all over the country. Some other pure exotic breeds such as Sirohi, Beetal, Jamnapari etc. and the crossbreds of BBG and exotic breeds are also available. According to Husain [1] and Amin et al. [2], more than 90% of the goats of the country are Black Bengal goat. BBGs are dwarf goats and are known to be famous for its adaptability, higher disease resistant, fertility, fecundity, early sexual maturity, larger litter size, delicacy of meat and superior skin quality, Devendra and Burns, [3]; Husain et al. [4]. Though majorities of the BBGs bear black coat color, but black and white, brown, brown and white, and white coat colors are also common in this population.

The phenotypic variation in a population arises due to genotypic and environmental effects, and the magnitude of phenotypic variability differs under different environmental conditions. Morphometric characters are continuous characters describing aspects of body shape, Dossa et al. [5]. Morphometric variation between populations can provide a basis for understanding flock structure, and may be more applicable for studying short-term, environmentally induced variation and thus more applicable to livestock management. According to Gizaw et al. [6], morphological description is an essential component of breed characterization that can be used to physically identify, describe, and recognize a breed, and also to classify livestock breeds into broad categories. Dossa et al. [5] reported that morphological measurements

such as heart girth, height at withers and body length can be used for rapid selection of large size individuals in the field to enable the establishment of elite flocks.

For animal selection purposes, it is essential to know how body shape of an animal behaves or in other word relationship among morphological measurements. It is also important to find out the relationship among different morphological traits for research purposes. However, body weight measurement at a regular interval is an important farm practice for domestic livestock. But frequently live body weight measurement is mostly a difficult task. Moreover, morphological body measurements with a measuring tape is less bothering than measuring live weight with a weighing scale. If we can estimate a regression equation considering body weight and body measurements it would be ease to estimate live body weight from a single or multiple body measurements like body length, heart girth, wither height etc. So, the present study was aimed to develop a regression model for estimating live body weight required with minimum morphometric information for BBG.

2. MATERIALS AND METHODS

2.1 Study Site

The research work was carried out at rural community level goat flocks at *Gangatia*, *Borachala* and *Pachpai* villages of Hobirbari Union under Bhaluka Upazila of Mymensingh district in Bangladesh from 2009 to 2013 under a completed project entitled "UNEP-GEF-ILRI FAnGR Asia Project".

2.2 Animals and Data

All the animals under this study were ear tagged to maintain individual identity as well as pedigree information. The morphometric measurements taken from 200 goats were collected from a baseline survey of different ages, sexes and colors along with 3 progressive generations produced from the foundation stock of the said community based goat breeding flocks. Separate data sheet for each animal was maintained for recording the information. Goat aged from birth to market age was used for phenotypic characterization and monthly data was recorded. All the flocks were maintained under the existing semi-intensive management system. Routine vaccination and de-worming against common diseases and parasites were conducted for all animals under the project.

2.3 Morphological Measurement and Analysis

Morphometric measurements of body length, chest girth, wither height and hip height along with body weights were recorded individually from birth up to 15 month age. All the information measured at a same age for sex and coat color were recorded. Body weights were measured in kilogram by a hanging spring balance and other morphological parameters were measured in centimeter by a measuring tape. The relationships amongst the morphometric body measures were determined using Pearson's

product moment correlation and linear regression procedure of SPSS 20.0 [7].

3. RESULTS AND DISCUSSION

The biometrical relationships between actual body weights and morphological body measurements according to sex are illustrated in Table 1. To develop regression equations for estimating live body weight from the external morphological measurements, four parameters viz. body length, chest girth, wither height and hip height were considered in this study. Thus, a total of seven regression models including single and multiple body measurements were considered. The analysis of variance showed that all the models were highly significant.

The regression analysis as illustrated in Table 1 shows that correlations between body weights with other morphological measurements are higher for female than that of male, except body length. The coefficients of determination were higher in female than that of male in the regression models including any of the single or multiple combinations of morphological traits. Tudu et al. [8] reported that coefficient of determination were 0.563 in male at 3 months, 0.529 in male at 9 months and 0.404 in female at 12 months ages, respectively. The relationships between body weight and morphological measurements according to coat color are shown in Table 2. The correlations and coefficients of determination were high and very close to each other, although slightly lower values are seen in

Table 1. Biometrical relationship between body measurement and body weight according to sex

Model information	Sex	n	R^2	Regression coefficients					Sig. level
				α	β_1	β_2	β_3	β_4	
Body length (BL)	Male	062	0.825	-13.008	0.598	-	-	-	***
	Female	138	0.811	-11.733	0.573	-	-	-	***
Chest girth (CG)	Male	062	0.842	-12.080	0.465	-	-	-	***
	Female	138	0.853	-10.737	0.445	-	-	-	***
Wither height (WH)	Male	062	0.782	-15.770	0.630	-	-	-	***
	Female	138	0.795	-14.688	0.621	-	-	-	***
Hip height (HH)	Male	062	0.799	-15.462	0.578	-	-	-	***
	Female	138	0.806	-13.990	0.558	-	-	-	***
BL+CG	Male	062	0.848	-12.696	0.203	0.313	-	-	***
	Female	138	0.856	-11.270	0.132	0.349	-	-	***
BL+CG+WH	Male	062	0.848	-12.601	0.193	0.306	0.022	-	***
	Female	138	0.857	-10.831	0.148	0.377	-0.059	-	***
BL+CG+WH+HH	Male	062	0.850	-13.082	0.187	0.279	-0.071	0.125	***
	Female	138	0.857	-10.848	0.148	0.371	-0.086	0.032	***

*n = number of observation; R^2 = coefficient of determination; α = constant value; β = regression coefficient; ***- $p < 0.001$

Table 2. Biometrical relationship between body measurement and body weight according to coat color

Model information	Coat color	n	R ²	Regression coefficients			Sig. level
				α	β ₁	β ₂	
Body length (BL)	Black	082	0.819	-12.141	0.580	-	***
	Black Bejoar	047	0.833	-11.482	0.549	-	***
	Brown Bejoar	110	0.812	-12.736	0.599	-	***
	Dutch belt	010	0.750	-12.034	0.555	-	***
Chest girth (CG)	Black	082	0.849	-11.163	0.452	-	***
	Black Bejoar	047	0.832	-10.674	0.423	-	***
	Brown Bejoar	110	0.852	-11.959	0.467	-	***
	Dutch belt	010	0.749	-9.696	0.413	-	***
BL+CG	Black	082	0.852	-11.652	0.141	0.348	***
	Black Bejoar	047	0.844	-11.348	0.285	0.209	***
	Brown Bejoar	110	0.858	-12.599	0.165	0.348	***
	Dutch belt	010	0.753	-11.050	0.305	0.188	***

* n = number of observation; R² = coefficient of determination; α = constant value; β = regression coefficient; ***-p<0.001

Table 3. Biometrical relationship between body measurement and body weight according to age

Model information	Age (month)	n	R ²	Regression coefficients			Sig. level
				α	β ₁	β ₂	
Body length (BL)	0 (Birth)	200	0.173	0.451	0.053	-	***
	1	200	0.266	1.195	0.067	-	***
	2	200	0.206	-0.067	0.147	-	***
	3	200	0.142	1.304	0.146	-	***
	6	200	0.393	-3.109	0.342	-	***
	9	200	0.305	-2.218	0.376	-	***
	12	200	0.374	-0.189	0.382	-	***
	15	200	0.225	6.507	0.271	-	***
Chest girth (CG)	0 (Birth)	200	0.106	0.564	0.040	-	**
	1	200	0.270	1.421	0.049	-	***
	2	200	0.111	1.623	0.078	-	***
	3	200	0.132	1.425	0.117	-	***
	6	200	0.388	-3.258	0.284	-	***
	9	200	0.242	-0.094	0.266	-	***
	12	200	0.402	-1.365	0.322	-	***
	15	200	0.485	-6.686	0.422	-	***
BL+CG	0 (Birth)	200	0.176	0.367	0.047	0.009	***
	1	200	0.319	1.083	0.043	0.023	***
	2	200	0.207	-0.161	0.137	0.011	***
	3	200	0.159	0.537	0.092	0.061	***
	6	200	0.484	-6.572	0.215	0.174	***
	9	200	0.345	-5.098	0.273	0.135	***
	12	200	0.492	-5.765	0.230	0.213	***
	15	200	0.487	-6.874	0.030	0.402	***

* n = number of observation; R² = regression coefficient; α = constant value; β = regression factors; ***-p<0.001; **-p<0.01; *-p<0.05)

Table 4. Biometrical relationship between body measurement and body weight irrespective of age, sex and coat color

Model information	R	R ²	Regression coefficients					Sig. level
			α	β_1	β_2	β_3	β_4	
Body length (BL)	0.904	0.817	-12.299	0.584	-	-	-	***
Chest girth (CG)	0.920	0.846	-11.321	0.453	-	-	-	***
Wither height (WH)	0.886	0.785	-15.041	0.621	-	-	-	***
Hip height (HH)	0.894	0.800	-14.565	0.565	-	-	-	***
BL+CG	0.922	0.851	-11.903	0.164	0.333	-	-	***
BL+CG+WH	0.922	0.851	-11.758	0.171	0.341	-0.020	-	***
BL+CG+WH+HH	0.923	0.851	-11.866	0.169	0.323	-0.090	0.089	***

*R = correlation coefficient; R² = coefficient of determination; α = constant value; β = regression coefficient

Table 5. Comparison between actual body weight and body weight estimated from regression equations developed by external body measurements according to age using model 5

Age at	Body weight in kg (Mean \pm SE)		Variation between actual and predicted		Correlation (R)
	Actual	Predicted	%	Sig. level	
Birth	1.50 \pm 0.02 (090)	1.51 \pm 0.01 (090)	0.67	NS	0.42**
1 month	3.03 \pm 0.04 (107)	3.02 \pm 0.02 (107)	0.30	NS	0.57**
2 month	4.67 \pm 0.09 (113)	4.69 \pm 0.04 (113)	0.43	NS	0.46**
3 month	6.61 \pm 0.11 (120)	6.59 \pm 0.04 (120)	0.30	NS	0.40**
6 month	10.94 \pm 0.17 (122)	10.96 \pm 0.12 (122)	0.18	NS	0.70**
9 month	14.50 \pm 0.22 (089)	14.47 \pm 0.13 (089)	0.21	NS	0.59**
12 month	17.89 \pm 0.28 (070)	17.87 \pm 0.19 (070)	0.11	NS	0.70**
15 month	19.68 \pm 0.34 (053)	19.69 \pm 0.24 (053)	0.05	NS	0.70**
Overall	8.77 \pm 0.22 (764)	8.79 \pm 0.20 (764)	0.23	NS	0.922**

NS-not significant ($p>0.05$); **-significant at 1% ($p<0.01$)

Table 6. Correlations between actual body weight and body weight estimated from regression equations developed by external body measurements

Model	Model information	Regression model	R
Model 2	Chest girth (CG)	$Y = \alpha + \beta \times CG$	0.920**
Model 3	Wither height (WH)	$Y = \alpha + \beta \times WH$	0.886**
Model 4	Hip height (HH)	$Y = \alpha + \beta \times HH$	0.894**
Model 5	BL+CG	$Y = \alpha + \beta_1 \times BL + \beta_2 \times CG$	0.922**
Model 6	BL+CG+WH	$Y = \alpha + \beta_1 \times BL + \beta_2 \times CG + \beta_3 \times WH$	0.922**
Model 7	BL+CG+WH+HH	$Y = \alpha + \beta_1 \times BL + \beta_2 \times CG + \beta_3 \times WH + \beta_4 \times HH$	0.923**

*R = correlation coefficient; α = constant value; β = regression coefficients; **- $p<0.01$)

Table 7. Correlations among the regression models

	Model 1	Model 2	Model 3	Model 4	Model 5	Model 6	Model 7
Model 1	1	0.962**	0.949**	0.948**	0.980**	0.980**	0.979**
Model 2		1	0.958**	0.964**	0.997**	0.997**	0.997**
Model 3			1	0.982**	0.963**	0.960**	0.960**
Model 4				1	0.967**	0.965**	0.969**
Model 5					1	1.000**	1.000**
Model 6						1	1.000**
Model 7							1

**significant at 1% level ($p<0.01$)

Dutch belt color. Tudu et al. [8] found that the coefficient of determination (R²) for height at wither and chest girth in combination (98.4%) was higher than other body measurements in

multiple traits evaluation for overall body weight of three color varieties of Bengal goats (Black, Brown and White) at 6 months of age.

The regression analyses according to different ages are given in Table 3 which shows that correlation coefficients as well as coefficients of determination increase as the age increases. The higher the ages the more accuracy of the prediction of body weight from morphological measurements. The coefficient of determination for combined tail length, body length and height at withers (86.1%) was higher than other body measurements in multiple traits evaluation for body weight in White Bengal goats at birth as reported by Tudu et al. [8]. They also found that the coefficient of determination (R^2) for height at withers (98.9%) and also for combined tail length, body length, height at withers and horn length (98.6%) were higher than other body measurements in multiple traits evaluation for body weight of Black Bengal goats at 9 and 12 months of age. Iqbal et al. [9] found that the coefficient of determination (R^2) for body length (64.8%) was higher than other body measurements in single trait evaluation in Beetal goat. They also reported that the coefficient of determination (R^2) was higher for height at withers and heart girth as compared to other body measurements.

The biometrical relationship between body weight and body measurement irrespective of sex, age coat color is illustrated in Table 4 which shows that correlation coefficients were higher when multiple body measurements were included in the model. Table 4 also shows that when chest girth as a single body measurement was included in the model gave the highest correlation coefficient. To get more accuracy, body length and chest girth are the best external body measures as coefficient of determination was high when included in the regression model. Tudu et al. [8] observed positive and significant correlation coefficients between body weight and linear body measurements with the exception of a few cases in all the three color varieties of Bengal goats. They also found that there was a high correlation between body weight and body length, height at withers and chest girth. Moela [10] found that the correlation coefficient between body weight and heart girth was high ($r = 0.91$), its direct effect on body weight was also the highest (path coefficient = 0.58) and significant ($P < 0.05$) as indicated by the t-test.

Table 5 shows the relationships between actual body weight and predicted body weight using

regression model 05 (using body length and chest girth) for different ages. In all ages, variations of body weight between actual and predicted were less than 1% ($p > 0.05$). The correlation coefficients for actual and predicted body weight were highly significant ($p < 0.01$) for all ages and magnitudes were higher in later ages. The overall correlation coefficient was 0.92** irrespective of age. Adeyinka and Mohammed [11] studied on Alpine ibex *Capra ibex* goat and observed that the squared value of the chest girth as predictive variables in the regression equation, being the most highly correlated linear measurements with total weight.

The correlations between actual body weight and body weight estimated from regression equations including single and multiple combination of external body measurements are presented in Table 6. The correlation coefficients for all models were significantly ($p < 0.01$) higher than 0.90 except model 3 and model 4. Table 6 also reveals that for a single morphological trait, chest girth is the best determinant for estimating live body weight and for multiple traits, body length along with chest girth are best determinant for estimating live body weight more accurately. Adeyinka and Mohammed [11] studied on Alpine ibex *Capra ibex* goat and they found the best prediction using age and different linear measurements in stepwise multiple regression.

Table 7 shows the relationships among regression models. The relationships among models were very strong and highly significant ($p < 0.01$). The correlation coefficients were very close to 1 which indicates that any of the models is suitable for estimating live body weight accurately. Iqbal et al. [9] reported the best fitted regression model in adult Beetal goats (13-18 months of age) to be body length and height at withers on body weight.

4. CONCLUSION

As a result of obtaining strong correlations among morphometric body measurements, body weight can be estimated from a single or multiple body measurements by establishing regression equations. However, for any single body measurement, chest girth is the best predictor and for multiple body measurements, body length along with chest girth is the best predictors for estimating live body weight more accurately.

ACKNOWLEDGEMENT

Authors would like to acknowledge “UNEP-GEF-ILRI FAnGR Asia Project” for providing fund and community goat keepers from where data were collected for this study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Husain SS. A study on the productive performance and genetic potentials of Black Bengal goats. A Ph.D. Thesis, Bangladesh Agricultural University, Mymensingh. 1993;3-108.
2. Amin MR, Husain SS, Islam ABMM. Reproductive peculiarities and litter weight in different genetic groups of Black Bengal does. Asian-Australasian Journal of Animal Science. 2001;14(3):297-301.
3. Devendra C, Burns M. Goats production in the tropics (Revised edn.). Tecl Comm., Bureaux of Animal Breeding and Genetics, Commonwealth Agricultural Bureaux, England. 1983;183.
4. Husain SS, Amin MR, Islam ABMM. Goat productions and its breeding strategy in Bangladesh. Proc. 1st National Workshop on Animal Breeding held on 26 Novenber, Bangladesh Agricultural University, Mymensingh, Bangladesh. 1998;17-36.
5. Dossa LH, Wollny C, Gauly M. Spatial variation in goat populations from Benin as revealed by multivariate analysis of morphological traits. Small Rum Res. 2007;73:150–159. [Cross Ref] DOI: 10.1016/j.smallrumres.2007.01.003
6. Gizaw S, van Arendonk JAM, Komen H, Windig JJ, Hanotte O. Population structure, genetic variation and morphological diversity in indigenous sheep of Ethiopia. Animal Genetics. 2007;38:621–628. [PubMed] [Cross Ref] DOI: 10.1111/j.1365-2052.2007.01659.x
7. SPSS Version 20.0. Statistical packages for Social Sciences, Version 20.0, SPSS Inc., Illinois, USA; 2008.
8. Tudu NK, Pyne SK, Ghosh N. Relationship of body weight with linear body measurements in three color varieties of Bengal Goats. International Journal of Scientific Research. 2015;4(2):16-20.
9. Iqbal M, Javed K, Ahmad N. Prediction of body weight through body measurements in Beetal goats. Journal of Animal and Plant Sciences. 2013;65(4):458-461.
10. Moela AK. Assessment of the relationship between body weight and body measurements in indigenous goats using path analysis. Master of Agricultural Management (Animal Production) Thesis. University of Limpopo, South Africa. 2014;1-56.
11. Adeyinka IA, Mohammed ID. Accuracy of body weight prediction in Nigerian Red Sokoto Goats Raised in North Eastern Nigeria using linear body measurement. Pakistan Journal of Biological Sciences. 2006;9(15):2828-2830.

© 2019 Habib et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/48413>